CarbonBrief
CLEAR ON CLIMATE

Factcheck: Trump’s climate report includes more than 100 false or misleading claims

By Ayesha Tandon, Leo Hickman, Cecilia Keating and Robert McSweeney
Design by Tom Prater

14 August 2025



Pages highlighted red contain false statements, whilst pages highlighted orange contain misleading statements. Pages can contain more than one false or misleading statement. Pages that remain uncoloured represent parts of the report that either have been stated as accurate by the cited author, or have not received any comment from invited experts. Front pages, reference and glossary pages omitted.

Executive summary

Page 0

MISLEADING

Elevated concentrations of CO_2 directly enhance plant growth, globally contributing to “greening” the planet and increasing agricultural productivity.

Dr David Lobell, associate professor, Stanford University.

I see two main problems with this (very old) argument. The direct benefits of CO_2 are widely acknowledged and nothing new. But we know that elevated CO_2 leads to climate changes and so the question is whether the CO_2 benefits are big enough to offset the climate losses. Their report does not address the net effects, which many studies have shown are negative, even for the US. The numbers they cite for direct effects of CO_2 are mainly from co2science.org, which is not a reputable source. Their summaries are not peer reviewed and include many studies of pots in greenhouses which are known to be biased. The numbers cited in the report are more than 2x what the best literature shows, such as in Ainsworth & Long (2021).

Quote from external source

Supporting evidence

1. Global Change Biology | Environmental Change Journal | Wiley Online Library

Page 0

FALSE

The world’s several dozen global climate models offer little guidance on how much the climate responds to elevated CO_2...The combination of overly sensitive models and implausible extreme scenarios for future emissions yields exaggerated projections of future warming.

Prof Veronika Eyring, professor of climate modelling, German Aerospace Center & University of Bremen.

In IPCC AR6, inconsistencies between models and observations and potential biases in model climate sensitivity values are taken into account in assessments of future climate change. IPCC AR6 clearly notes that some CMIP6 models simulate warming that is either above or below the assessed very likely range of observed warming. However, this information about how well models simulate past warming, as well as other insights from observations and theory, are used to assess projections of global warming (see cross-section Box TS.1 in Arias et al. (2021)).

Quote given to Carbon Brief

Supporting evidence

1. IPCC AR6 Working Group 1: Technical Summary | Climate Change 2021: The Physical Science Basis

Part I: Direct human influence on ecosystems and the climate

1 Carbon dioxide as a pollutant

Page 2

MISLEADING

The growing amount of CO_2 in the atmosphere directly influences the Earth system by promoting plant growth (global greening), thereby enhancing agricultural yields, and by neutralising ocean alkalinity.

Anonymous.

This ignores other effects of rising CO_2 concentrations, i.e.: on climate. It is also failing to mention that increased CO_2 can reduce the nutrient density of some crops.

Quote given to Carbon Brief

Supporting evidence

1. Chapter 5: Food, Fibre and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.1 CO2 as a contributor to global greening

Page 3

FALSE

While plant models predict increased photosynthesis in response to rising CO_2, Haverd et al. (2020) reported a CO_2 fertilisation rate much larger than model predictions. That is, CO_2 fertilisation had driven an increase in observed global photosynthesis by 30% since 1900, versus 17% predicted by plant models. If true it would indicate that global models of the socioeconomic impacts of rising CO_2 have understated the benefits to crops and agriculture.

Dr Delphine Deryng, lead author, IPCC AR6 WG2.

The paper by Haverd et al. focuses on natural ecosystems, not crops. So whilst the findings that CO_2 fertilisation effects on global greening makes a larger share relative to other factors, the results are not directly transferable to the socio-economic impacts of rising CO_2 on agriculture. Rising CO_2 contributes to higher radiative forcing which increases global mean temperature and accelerates the global water cycle, causing increases in the severity and frequency of extreme weather events (e.g. droughts, heat-stress and wildfires), particularly threatening crop yields and production.

Quote given to Carbon Brief

Supporting evidence

1. Global Change Biology | Environmental Change Journal | Wiley Online Library

2.1 CO2 as a contributor to global greening

Page 3

MISLEADING

The growing CO_2 concentration in the atmosphere has the important positive effect of promoting plant growth by enhancing photosynthesis and improving water use efficiency.

Anonymous.

Promoting plant growth is not always positive, because some species benefit more than others, which creates risks to biodiversity. For example, in tropical forests, elevated CO_2 promotes the growth of lianas, which are parasites that threaten trees. Also increased CO_2 fertilisation is playing a role in disrupting grassland and savannah ecosystems by promoting tree and shrub growth ("woody encroachment").

Quote given to Carbon Brief

Supporting evidence

1. Increasing dominance of large lianas in Amazonian forests | Nature
2. Chapter 2: Terrestrial and Freshwater Ecosystems and Their Services | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.1 CO2 as a contributor to global greening

Page 3

FALSE

Section 2.1.1

Anonymous.

The following section cites papers that are, in fact, showing evidence for other drivers of change, e.g. page 14 line 34 says "Piao et al. (2020) noted that greening was even observable in the Arctic", but Piao et al (2020) actually show that warming is the dominant driver of greening in the Arctic, not CO_2 fertilisation (see figure 4 and associated text). Also the authors of the DoE report contradict their own statement two paragraphs later by saying: "Chen et al. (2019) show that in China and India much of it is driven by land management changes."

Quote given to Carbon Brief

Supporting evidence

1. Characteristics, drivers and feedbacks of global greening | Nature Reviews Earth & Environment

2 Direct impacts of CO2 on the environment

Page 3

MISLEADING

Piao et al. (2020) and Chen et al. (2024) report that the greening trend continues with no evidence of slowdown.

Anonymous.

The DoE authors fail to mention another study which shows the opposite, that greening was reversed around the year 2000 over 90% of the global vegetated area.

Quote given to Carbon Brief

Supporting evidence

1. Inhibitive Effects of Recent Exceeding Air Temperature Optima of Vegetation Productivity and Increasing Water Limitation on Photosynthesis Reversed Global Greening - Chen - 2022 - Earth's Future - Wiley Online Library

2.1 CO2 as a contributor to global greening

Page 3

MISLEADING

Section 2.1

Dr David Crisp, retired atmospheric physicist.

Chapters 2 and 9 assert that CO_2 fertilisation will increase plant growth and crop yields. The proposed benefits of CO_2 fertilisation are not realised in this set of DGVMs because this is only one of several mechanisms that control plant growth. As any farmer knows, plant growth is rarely limited by the abundance of the most abundant nutrient. It is usually limited by the abundance of the least abundant nutrient. While increased CO_2 can accelerate plant growth in carefully controlled laboratory conditions, this rarely happens in nature or in large-scale agriculture. There, plant growth is usually limited by water, nitrogen, phosphorus, sunlight or temperature. These models include all of those effects. The range of outputs produced by the models reflects uncertainties in the relative roles of these processes and their potential evolution with climate change. This behaviour should foster serious concern (doubt) about the potential benefits of CO_2 fertilisation in a changing climate. There is no discussion of this here or in chapters 2.1 or 9.

Quote given to Carbon Brief

Supporting evidence

2.1 CO2 as a contributor to global greening

Page 4

MISLEADING

Had atmospheric CO_2 levels continued declining, plant growth would have declined and eventually ceased. Below 180ppm, the growth rates of many C3 species are reduced 40-60% relative to 350ppm (Gerhart and Ward (2010)) and growth has stopped altogether under experimental conditions of 60-140ppm CO_2. Some C4 plants are still able to grow at levels even as low as 10ppm, albeit very slowly (Gerhart and Ward (2010)).

Prof Joy Ward, provost and executive vice president, Case Western Reserve University.

“Ward, however, told WIRED in an emailed statement that her experiments were conducted under ‘highly controlled growth conditions’ to create a ‘mechanistic understanding’ of CO_2, and that climate change can cause a host of impacts on plants not accounted for in her study. ‘With rising CO_2 in natural ecosystems, plants may experience higher heat loads, extreme weather events such as droughts and floods and reduced pollinators – which can have severe net negative effects on plant growth and crop yields,’ she says. ‘Furthermore, our studies indicate that major disruptions in plant development such as flowering time can occur in direct response to rising CO_2, which were not mentioned in the report.’”

Quote from external source

Supporting evidence

2.1 CO2 as a contributor to global greening

Page 4

FALSE

Piao et al. (2020) and Chen et al. (2024) report that the greening trend continues with no evidence of slowdown, and CO_2 fertilisation remains the dominant driver.

Dr Delphine Deryng, lead author, IPCC AR6 WG2.

Chen et al (2024) does not back up this statement. In the abstract, the author concluded that greening, whilst still increasing, has slowed down: "Our study highlighted that drought trend did not necessarily trigger vegetation browning, but slowed down the rate of greening." Piao et al. (2019) looked at the driver of global greening. They did not look at whether trends in global greening are rising or decreasing. In addition they analysis focused on historical observation (1980-2010) and did not assess future trends.

Quote given to Carbon Brief

Supporting evidence

1. The global greening continues despite increased drought stress since 2000 - ScienceDirect
2. Characteristics, drivers and feedbacks of global greening | Nature Reviews Earth & Environment

2.1 CO2 as a contributor to global greening

Page 4

MISLEADING

Had atmospheric CO_2 levels continued declining, plant growth would have declined and eventually ceased. Below 180ppm, the growth rates of many C3 species are reduced 40-60% relative to 350ppm (Gerhart and Ward (2010)) and growth has stopped altogether under experimental conditions of 60-140ppm CO_2. Some C4 plants are still able to grow at levels even as low as 10ppm, albeit very slowly (Gerhart and Ward (2010)).

Anonymous.

The decline in atmospheric CO_2 levels over the last few tens of millions of years stopped naturally, and for the last 800,000 years up until the Industrial Revolution did not show much of a trend, just fluctuating between about 170 and 280 parts per million. The hypothetical scenario of a further decline below these levels is not relevant – it is not the case that human-driven CO_2 emissions have somehow saved us from declining CO_2 levels and declining plant growth, as seems to be the implication behind this paragraph.

Quote given to Carbon Brief

Supporting evidence

1. Chapter 5: Global Carbon and other Biogeochemical Cycles and Feedbacks | Climate Change 2021: The Physical Science Basis

2.1 CO2 as a contributor to global greening

Page 5

MISLEADING

The overwhelming theme is that plants, especially C3 plants, benefit from extra CO_2.

Anonymous.

While individual plants benefit in isolation, the overall effect on an ecosystem and biodiversity can be detrimental due to some species benefitting more than others and out-competing them – for example, lianas responding more than trees, which they damage, encroachment of trees and shrubs into grasslands and savannahs, and the promotion of invasive species and weeds.

Quote given to Carbon Brief

Supporting evidence

1. Increasing dominance of large lianas in Amazonian forests | Nature
2. Chapter 2: Terrestrial and Freshwater Ecosystems and Their Services | Climate Change 2022: Impacts, Adaptation and Vulnerability
3. Chapter 5: Food, Fibre and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.1 CO2 as a contributor to global greening

Page 5

MISLEADING

There are two mechanisms by which CO_2 confers a growth benefit

Anonymous.

A "growth benefit" to the plant is not necessarily beneficial in other ways – for example, from IPCC AR6: "Perennial crops and root crops may have a greater capacity for enhanced biomass under elevated CO_2 concentrations, although this does not always result in higher yields. For some food crops, nutrient density declines due to elevated CO_2." And: "Elevated CO_2 reduces some important nutrients such as protein, iron, zinc and some grains, fruit or vegetables to varying degrees depending on crop species and cultivars."

Quote given to Carbon Brief

Supporting evidence

1. Chapter 5: Food, Fibre and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.1 CO2 as a contributor to global greening

Page 5

MISLEADING

The gains induced by increasing CO_2 from 150ppm to 350ppm continue under a further doubling to 700ppm.

Anonymous.

This very simplistic illustration from a small laboratory study ignores key effects such as nutrient availability, which in the real world can constrain the response to elevated CO_2.

Quote given to Carbon Brief

Supporting evidence

1. Global Change Biology | Environmental Change Journal | Wiley Online Library

2.2 The alkaline oceans

Page 6

MISLEADING

Section 2.2.1

Dr John Kennedy, climate scientist.

"The first subsection builds towards the conclusion in the summary that ‘ocean life is complex and much of it evolved when the oceans were acidic relative to the present’. Leaving aside the vacuity of this argument for the moment – life itself evolved when there was little oxygen in the atmosphere, the biochemical innovations that flooded the atmosphere with oxygen were catastrophic for life then, but see how we would do without it now – their citations are spare and strange particularly within the context of later arguments about the value of models and the contention that this is some sort of meaningful ‘critical review’. The first paper they cite regarding long-term change in ocean pH is Krissansen-Totton et al. (2018), which uses a model to constrain climate and ocean pH of the early Earth up to the present. They find that ocean pH evolves monotonically from 6.6 (see the abstract of the paper for the broad uncertainty ranges) at 4.0 Ga to 7.0 at the Archean-Proterozoic boundary, and to 7.9 at the Proterozoic-Phanerozoic boundary reaching a modern value of 8.2. While we might raise an eyebrow that the ‘critical review’ finds models are good enough for the herculean task of reproducing almost the whole of Earth’s climate history, but not for understanding the past 200 years, the eyebrow is likely to go shooting off your face when you reach the sentence in the ‘critical review’ that says: ‘Even if the water were to turn acidic, it is believed that life in the oceans evolved when the oceans were mildly acidic with pH 6.5 to 7.0.’ I derive little comfort from the fact that simple life forms evolved in such conditions. The gist of the ‘critical review’ isn’t that simple life forms will survive the current warming, but that human society supported by a flourishing biosphere will not just survive but thrive. Anyway, this is only part of an argument and the whole of the argument is never really spelled out. It seems to go something like this: pH of the ocean varied in the past and we exist today, therefore we will always exist and pH of the ocean is unimportant.

"The second, shorter, long-term perspective, which feeds into this argument, mixes up surface pH (as shown in Figure 2.3 from the CMEMS dataset) with deep ocean pH (from Rae et al. (2018)) who (according to their abstract) ‘present deep-sea coral boron isotope data that track the pH – and thus the CO_2 chemistry – of the deep Southern Ocean over the past forty thousand years’. ‘Deep’ and ‘track’ are the operative words here. The estimated changes are ‘deep’, from a depth of around 750 metres and not the surface. Regarding ‘track’, the numbers quoted in the ‘critical review’ – pH of 7.4 to 7.5 20,000 years ago – came presumably from Figure S1 in the Rae paper, which provides an approximate conversion of the boron isotopes to pH. How very very approximate they are is shown by an inset uncertainty range, which extends from well below 7.4 to well above 7.6 suggesting great care is needed in the interpretation of the absolute pH values."

Quote from external source

Supporting evidence

1. Constraining the climate and ocean pH of the early Earth with a geological carbon cycle model | PNAS

2.2 The elusive oceans

Page 5

MISLEADING

Human CO2

Dr John Kennedy, climate scientist

> Dr John Kennedy: [...] The oceans are the great buffer for the climate system. Without the oceans, the atmospheric carbon dioxide concentration would be rising much faster than it is at present. The oceans have taken up around a quarter of the carbon dioxide that we've released into the atmosphere since the start of the industrial revolution. Another way of looking at it is that we've released around 500 billion tonnes of carbon dioxide into the atmosphere since the start of the industrial revolution, and around 125 billion tonnes of that is now in the oceans.

Anonymous,

> The rest of the section concerns itself with the more general impacts of ocean acidification. And with good reason. Only the Devereux (2016) on the tails of reef results in the literature and offer, as an example, Clements et al. (2017) which is about the direct effects of CO2 on the behaviour of fish specifically (not the reefs themselves) though it does have a risky metanalysis component that serves little purpose of suggesting that discussion of the topic is one-sided. If anything, Clements et al. show that the ocean acidification issue is not as bad as it is made out to be in the popular media. The Devereux article is also somewhat meta and points out that papers on ocean acidification overwhelmingly show that CO2 is bad for the ocean and that only 43% and 36% of studies with p<0.05 are in the correct direction for pH and CO2 levels that are seen there now!! I don't know. The Devereux and Clements articles are both still on a far extreme of a strong bias. A simple Google Scholar search will find you that not only are there huge numbers of papers mentioning the topic in the past five years, but there are even lists of review papers and meta-analyses on the topic which cover a much broader range of impacts. Summarizing that literature with just 12 references (Devereux) or 15 (Clements) in order to suit adequate my my definition.

Quote from second section

Supporting evidence

- Chandler, J.C. Australasian, J. Climate Research 2022: Impacts, Adaptation and Vulnerability
- A subsequent section looks to be very helpful on the topic (Kroeker et al. Global Change Biology - Ocean Acidification and Warming Impacts)
- Meta-analysis of ocean acidification responses to ocean acidification research: ICES Journal of Marine Science (Oxford Academic)
- Meta-analysis reveals an extreme "decline effect" in the impacts of ocean acidification on fish behavior | PLOS Biology

Quote from [second section](#)

2.2.1 CO2 as a contributor to global warming

Page 5

MISLEADING

Denying et al. (2016) summarised evidence on crop water productivity of 78%, the yield gain of well-water used, showing attention to the quantification of CO2 effects on crop water productivity. Experiments and re-analyses local and cosmographic master lines during real experiments. They surveyed all available FACE (free air carbon dioxide enrichment) studies and found that crop yield changes for maize, wheat, rice, and soybean and combined it with crop models data simulating yield responses as a function of the extreme CO2 concentrations. Their results, in four growing regions (tropics, and, temperate and cold) each of which were split into irrigated and rainfed conditions, showed that crop water productivity increased significantly for all four crops in all regions. They were more than offset by CO2 fertilization, as the net regional responses of yield to CO2 ranged from 11% to 25% and those of water use efficiency (WUE) from 26% to 41% (Table 1). Their results are evidence that the global models underestimate the CO2 fertilization effect.

Dr Delphine Deryng, lead author, IPCC WG1 WG2

> Dr Delphine Deryng: [...] Our study is the first to look at how climate change impacts the interannual variation of global crop yields. Using statistical crop models, focusing specifically on how water affects crop yield and evapotranspiration, we found that climate change has already had a negative impact on global crop production. We also found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past. Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. We found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past.

Quote from [Nature Food](#)

Supporting evidence

- [FAO](#)
- Overcoming uncertainties in the effects of elevated CO2 on crops | Nature Food
- Multimodel assessment of water scarcity under climate change | PNAS

2.2.1 CO2 as a contributor to global warming

Page 5

FALSE

Denying et al. (2016) assessed that climate change would "exacerbate water scarcity". [...] The results do predict that global crop water productivity will increase significantly. Given that the agronomic greening is happening even in arid areas

Dr Delphine Deryng, lead author, IPCC WG1 WG2

> Dr Delphine Deryng: [...] Our study is the first to look at how climate change impacts the interannual variation of global crop yields. Using statistical crop models, focusing specifically on how water affects crop yield and evapotranspiration, we found that climate change has already had a negative impact on global crop production. We also found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past. Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. We found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past.

Quote from [Nature Food](#)

Supporting evidence

- Global forces in human water security and river biodiversity | Nature
- Multimodel assessment of water scarcity under climate change | PNAS

2.2.2 CO2 as a contributor to global warming

Page 5

FALSE

The IPCC has only reasonably discussed global greening and CO2 fertilization of agricultural crops. The topic is briefly acknowledged in a few places in the body of the AR5 text and most extensively treated in a statistical sub-section of Section 2.2.4.3 of the 600+ WG1 report, "global greening and browning," points out that [...] the IPCC doesn't even cite Chandler (2022) which is the most recent publication that has greening had increased globally over the past two to three decades. It then discusses the three new estimates in the greening and browning data sets, concluding that while they have high confidence greening has occurred, they have low confidence in the magnitude of the trend. There are also brief mentions in Section 5.4.8.1 of elevated CO2 as both a net benefit and a disbenefit to crop yields (but not WUE) and 5 synthesis

Dr Delphine Deryng, lead author, IPCC WG1 WG2

> Dr Delphine Deryng: [...] Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. Using statistical crop models, focusing specifically on how water affects crop yield and evapotranspiration, we found that climate change has already had a negative impact on global crop production. We also found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past. Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. We found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past.

Quote from [Nature Food](#)

Supporting evidence

- Chandler J. Fisher and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.2.2 CO2 as a contributor to global warming

Page 5

FALSE

Overall, however, the polycrystalline nature, technical assessment and synthesis reports of AR5 and AR6 do not change the conclusions.

Dr Delphine Deryng, lead author, IPCC WG1 WG2

> Dr Delphine Deryng: [...] Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. Using statistical crop models, focusing specifically on how water affects crop yield and evapotranspiration, we found that climate change has already had a negative impact on global crop production. We also found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past. Our study is the first study to look at how climate change impacts the interannual variation of global crop yields. We found that irrigation can help to reduce the negative impacts of climate change on crop yields. Irrigation can also help to reduce the variability in crop yields that we've observed in the past.

Quote from [Nature Food](#)

Supporting evidence

- Technical Summary | Climate Change 2022: Impacts, Adaptation and Vulnerability
- Summary for Policymakers | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.2.2 CO2 as a contributor to global warming

Page 5

MISLEADING

The IPCC has only reasonably discussed global greening and CO2 fertilization of agricultural crops.

Anonymous,

> The IPCC has extensively discussed the protection of plant growth by rising atmospheric CO2 concentrations and consequent effects on plant growth and water use efficiency. The topic is briefly acknowledged in a few places in the body of the AR5 text and most extensively treated in a statistical sub-section of Section 2.2.4.3 of the 600+ WG1 report, "global greening and browning," points out that [...] the IPCC doesn't even cite Chandler (2022) which is the most recent publication that has greening had increased globally over the past two to three decades. It then discusses the three new estimates in the greening and browning data sets, concluding that while they have high confidence greening has occurred, they have low confidence in the magnitude of the trend. There are also brief mentions in Section 5.4.8.1 of elevated CO2 as both a net benefit and a disbenefit to crop yields (but not WUE) and 5 synthesis

Quote from [second section](#)

Supporting evidence

- Chandler J. Fisher and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.2.2 CO2 as a contributor to global warming

Page 5

MISLEADING

The IPCC has only reasonably discussed global greening and CO2 fertilization of agricultural crops.

Anonymous,

> The IPCC has extensively discussed the protection of plant growth by rising atmospheric CO2 concentrations and consequent effects on plant growth and water use efficiency. The topic is briefly acknowledged in a few places in the body of the AR5 text and most extensively treated in a statistical sub-section of Section 2.2.4.3 of the 600+ WG1 report, "global greening and browning," points out that [...] the IPCC doesn't even cite Chandler (2022) which is the most recent publication that has greening had increased globally over the past two to three decades. It then discusses the three new estimates in the greening and browning data sets, concluding that while they have high confidence greening has occurred, they have low confidence in the magnitude of the trend. There are also brief mentions in Section 5.4.8.1 of elevated CO2 as both a net benefit and a disbenefit to crop yields (but not WUE) and 5 synthesis

Quote from [second section](#)

Supporting evidence

- Chandler J. Fisher and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.2.2 CO2 as a contributor to global warming

Page 5

FALSE

Overall, however, the polycrystalline nature, technical assessment and synthesis reports of AR5 and AR6 do not change the conclusions.

Anonymous,

> The topic of CO2 fertilization effects, including their limitations and interactions with other drivers of change, are extensively discussed in AR5 WGII Chapter 5, Food, Fibre, and Other Ecosystem Products. The term 'CO2' appears 156 times in that chapter. In the WG1 report, Chapter 5 is 'Global Carbon and Other Biogeochemical Cycles and Feedbacks' and, again, extensively includes discussion of CO2 effects as part of the carbon cycle.

Quote from [second section](#)

Supporting evidence

- Chandler J. Fisher and Other Ecosystem Products | Climate Change 2022: Impacts, Adaptation and Vulnerability

2.2.3 CO2 as a contributor to global warming

Page 5

MISLEADING

Even if the ocean were to turn acidic, it is believed that life in the oceans evolved when the oceans were mildly acidic with pH 3 to 7.8 (Hönisch+ Tomas et al. (2012).

Dr Jamie Kaas, marine biochemistry and climate scientist

> Modern marine life and ecosystems e.g. coral reefs and open ocean fisheries are very different from the initial fisheries that evolved in the earliest oceans e.g. bacteria, single-celled organisms and so have different adaptive ranges and vulnerabilities. In the early oceans the optimal temperature conditions which were suitable for diseases have little relevance for optimal conditions for humans, there is little relevance in comparing the optimal natural habitats to the conditions in which modern marine life has evolved.

Quote from [Carbon Brief](#)

Supporting evidence

2.2.3 CO2 as a contributor to global warming

Page 5

MISLEADING

Even if the ocean were to turn acidic, it is believed that life in the oceans evolved when the oceans were mildly acidic with pH 3 to 7.8 (Hönisch+ Tomas et al. (2012).

Dr Jamie Kaas, marine biochemistry and climate scientist

> Modern marine life and ecosystems e.g. coral reefs and open ocean fisheries are very different from the initial fisheries that evolved in the earliest oceans e.g. bacteria, single-celled organisms and so have different adaptive ranges and vulnerabilities. In the early oceans the optimal temperature conditions which were suitable for diseases have little relevance for optimal conditions for humans, there is little relevance in comparing the optimal natural habitats to the conditions in which modern marine life has evolved.

Quote from [Carbon Brief](#)

Supporting evidence

2.2.3 CO2 as a contributor to global warming

Page 5

MISLEADING

On the time scales of thousands of years, boron isotope proxy measurements show that ocean pH was around 7.8 or 7.9 during the last glaciation (up to about 30,000 years ago) increasing to present day values as the world warmed during deglaciation (Hat et al. 2012).

Dr Jamie Kaas, marine biochemistry and climate scientist

> Modern marine life and ecosystems e.g. coral reefs and open ocean fisheries are very different from the initial fisheries that evolved in the earliest oceans e.g. bacteria, single-celled organisms and so have different adaptive ranges and vulnerabilities. In the early oceans the optimal temperature conditions which were suitable for diseases have little relevance for optimal conditions for humans, there is little relevance in comparing the optimal natural habitats to the conditions in which modern marine life has evolved.

Quote from [Carbon Brief](#)

Supporting evidence

- Atmospheric Ocean CO2 Exchange Across the Last Deglaciation Predicted by the Boron Isotope Proxy - Hain - 2021 - Paleoceanography and Paleoclimatology - Wiley Online Library
- Last Glacial and Holocene Ocean pH: New Constraints on Geochemical Interpretation | Science

2.2.4 CO2 as a contributor to global warming

Page 5

MISLEADING

Thus, ocean biota appear to be resilient to natural long-term changes in ocean pH when marine organisms were exposed to such ranges in pH.

Dr Jamie Kaas, reader, geochemistry and climate, University of St Andrews

> Natural long-term changes in ocean pH have a fundamentally different impact on ocean chemistry than rapid impact of ocean chemistry from fossil fuels (see Hönisch et al. (2012). When ocean pH is lowered rapidly (e.g. on decadal-to-centennial timescales), the abundance of carbonate ions decreases rapidly. As these carbonate ions are the most critical ingredient for marine shell formation, shells become harder to grow. Historically, this is known as a key mechanism responsible for mass extinction and it makes calcium carbonate shell formation more energetically costly (e.g. Gagnon et al., (2021). In contrast, when ocean pH is lowered slowly, as it is on timescales of a thousand years or more, the dissolution of seafloor calcium carbonate, alongside weathering of rocks on land, can act to counteract the change in ocean pH. This is because the dissolution of carbonate sediments shapes the lower pH. Today, anthropogenic CO2 emissions is lowering ocean pH on decadal-to-centennial timescales, which are too rapid for these natural processes to help us, and leading to long-term calcium carbonate saturation states, and making shell formation more energetically costly and enhancing environmental stress on these organisms.

Quote from [Carbon Brief](#)

Supporting evidence

- The Geological Record of Ocean Acidification: Science
- Controls on Ocean Acidification Its Past, Present, and Future Modulation | Annual Reviews

2.2.4 CO2 as a contributor to global warming

Page 5

MISLEADING

Similarly, a meta-analysis (Clements et al. (2021) of the negative effects of ocean acidification on reef fish behaviour finds that the majority of studies (91%) with "no effect" (i.e. natural) dramatic conclusions published in journals showing apparently large impacts. A publication related to the followed up by subsequent publications with larger sample sizes finding much smaller and typically non-existent effects. This call for their colleagues to improve research practices to counter the "decline effect" ("the tendency of initially exciting, large effect sizes in this field tend to be characterized by low sample size, as yet unpublished, and highly variable outcomes") and recommended that publication bias be addressed in all meta-analyses. No comment that ocean acidification may have an overestimated effect on fish behaviour, and we advocate for improved approaches to minimize the potential for the decline effect to distort science (Clements et al. (2021).

Dr Jeff Clements, marine ecologist, Fisheries and Oceans Canada (Gulf Region)

> "Clements... says that the way the Gulf report cites his research on ocean acidification and fish behavior is accurate. "From an empirical standpoint", Clements says in an email to Carbon Brief, "I would argue that there is no evidence that ocean acidification is a threat to marine ecosystems and does not mean that a myriad of other ocean ecosystems, biological processes and species will fare similarly. Other work from his lab, meanwhile, has underscored the vulnerability of mussels to ocean warming and looked at how boat wakes negatively affect clam behavior... "I want to make it clear that our research does not suggest that ocean acidification is not a problem, but it may not be more generally is not a problem," he tells WIRED. "While effects on fish behavior may, or may not, be an ocean acidification issue, species and biological processes are certainly vulnerable to the impacts of acidification and the compendium of other climate change stressors that our oceans are experiencing."

Quote from [original source](#)

Supporting evidence

2.2.4 CO2 as a contributor to global warming

Page 5

MISLEADING

Similarly, a meta-analysis (Clements et al. (2021) of the negative effects of ocean acidification on reef fish behaviour finds that the majority of studies (91%) with "no effect" (i.e. natural) dramatic conclusions published in journals showing apparently large impacts. A publication related to the followed up by subsequent publications with larger sample sizes finding much smaller and typically non-existent effects.

Prof Fredrik Ladefoged, professor of Pre-ecophysiology, Norwegian University of Science and Technology

The way the IAM report cites our own work is flawed, as they use a specific finding in a broad subsection of the literature to support claims that are not in line with it. This is intellectually dishonest and unscientific, and suggests either gross incompetence or an agenda to mislead.

Quote given to Carbon Brief

Supporting evidence

1. [Were analytics trends an extreme "decline effect" in the impacts of ocean acidification on fish behavior?](#) - Nick Raby

2.2 The alkaline oceans

Page 5

MISLEADING

In summary, ocean life is complex and much of it evolved when the oceans were acidic relative to the present. The ancestors of modern coral first appeared about 245m years ago, 225 levels far more than 280m years ago, and modern coral reefs more than 60m years back. Much of the public discussion of the effects of ocean "acidification" on marine biota has been one-sided and misleading.

Quote from [IAM report](#), [section 2.2.2.2](#)

Supporting evidence

2.2 The alkaline oceans

Page 5

MISLEADING

In summary, ocean life is complex and much of it evolved when the oceans were acidic relative to the present. The ancestors of modern coral first appeared about 245m years ago, 225 levels far more than 280m years ahead were many times higher than they are today. Much of the public discussion of the effects of ocean "acidification" on marine biota has been one-sided and misleading.

Dr James Rae, reader, geochemistry and climate, University of St Andrews.

As discussed above, long-term geological pH changes are not a good analogue for a rapid drop in pH of the type happening today, for the key reason that a rapid pH drop results in lowered calcium carbonate saturation state, which makes shell formation harder (e.g. Hönisch et al., 2012). To find better analogues for modern pH declines in the past, we need to look at periods in Earth's history when calcium carbonate saturation was low, such as the Paleocene-Eocene Thermal Maximum (e.g. Penman et al., 2014). These show that all of the major mass extinction events in the ocean for which we have pH reconstructions were associated with low pH oceans. Overall, the geological record does suggests that rapid pH decline is a barrier to serious concern for marine ecosystems and the communities and economies which they support.

Quote given to Carbon Brief

Supporting evidence

1. [The Carbon Record of Ocean Acidification: Science](#)
2. [Pulses of ocean acidification at the Triassic–Jurassic boundary? | Nature Communications](#)

References

Page 9

FALSE

Wang, Y., Hsieh, W.C.H., Chen Peter, Z. (2022). *The global planting potential to mitigate climate change through stomata since 1982. Global Ecology and Conservation, 40, e02356.*

Anonymous.

The author list incorrect on this citation. It should be Xiao Chen, Tian Chen, Bin Xu, Jian Zhu, Shengbiao Zhou and Yingjie Wang. It's details associates the paper with well-known authors and demonstrates lack of attention to detail.

Quote given to Carbon Brief

Supporting evidence

1. [The global potential to remove drastic increased drought stress since 1982 - ScienceDirect](#)

3 Human influences on the climate

Page 11

MISLEADING

The IPCC has downplayed the role of the sun in climate change, but there are significant solar irradiance reconstructions that imply it contributed to recent warming.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research

This is a vague and potentially misleading statement. The IPCC AR6 adopted the solar irradiance forcing values recommended by Matthes et al. (2016). Of the CMIP6 models in both cases, the TSI and solar irradiance reconstructions selected were the most commonly used and recommended ones. The selection of the spectral TSI and SSI variation that these were 'dependent' while assigning a low TSI, without examining in further detail the impact that these 'dependencies' have on the climate response. Additionally, the choice of the solar irradiance reconstructions implying a low variability was not to be considered. The most scientifically robust irradiance reconstructions indicate that this contribution has been relatively small in the order of 0.1°C since 1850 and that the apparent impacts to implicitly reference the work of Connolly et al. (2021) and (2023), although only the former is cited in the report. Connolly et al. (2021) argued that a significant portion, or even most, of recent global warming could be attributed to solar irradiance changes, which is inconsistent with the vast majority of solar irradiance reconstructions, right with low long-term variability and right with high variability. To resolve this balance, they conducted a sensitivity analysis of models with strongly varying low-variability reconstructions.

This is not correct and was tested by Connolly et al. (2021):

- Hoyt and Schatten (1993) has been thoroughly discredited (see Christoudias (2016)). It includes arbitrary scaling parameters and even fabricated data, with some cycles seemingly copied from another one's data.
- Lean et al. (1995) is an early version of the NRL (Naval Research Laboratory) model that has since been superseded by more advanced models (Coddington et al. (2016), Yeo, Krivova, Coddington et al. (2014), and Yeo et al. (2017)).
- Reid et al. (2000) used a simplified linear conversion from cosmogenic isotope production (e.g., 10Be to TSI, without considering the nonlinear physics in the geomagnetic modulation, atmospheric transport processes, or climate-driven reservoir changes. Reid and Turney (2008) used a similar simplified approach. Both studies lack STOKE-TI (Spectral And Total Irradiance Reconstructions) (Yeo et al. (2014)), which show significantly lower long-term variability. Furthermore, Steinhilber and Beer (2013) revised the original Reid et al. (2000) reconstruction which was further updated in Steinhilber et al. (2009). Both versions were tested in Connolly et al. (2021) and are included in the ensemble.
- Shapiro et al. (2011), an older high-variability model, was effectively replaced by Egorova et al. (2018), which itself has been criticised by Yeo et al. (2020) for exaggerating long-term TSI changes. Egorova et al. (2018) was included five times in the Connolly et al. (2021) ensemble, including Shapiro et al. (2011).
- Reid et al. (2000) used a simplified linear conversion from cosmogenic isotope production (e.g., 10Be to TSI, without considering the nonlinear physics in the geomagnetic modulation, atmospheric transport processes, or climate-driven reservoir changes. Reid and Turney (2008) used a similar simplified approach. Both studies lack STOKE-TI (Spectral And Total Irradiance Reconstructions) (Yeo et al. (2014)), which show significantly lower long-term variability. Furthermore, Steinhilber and Beer (2013) revised the original Reid et al. (2000) reconstruction which was further updated in Steinhilber et al. (2009). Both versions were tested in Connolly et al. (2021) and are included in the ensemble.
- Shapiro et al. (2011), an older high-variability model, was effectively replaced by Egorova et al. (2018), which itself has been criticised by Yeo et al. (2020) for exaggerating long-term TSI changes. Egorova et al. (2018) was included five times in the Connolly et al. (2021) ensemble, including Shapiro et al. (2011).

Crucially, in all these high-variability models, the long-term TSI trends were not derived from observational constraints but were imposed as assumptions. Modern versions of the NRL model (Lean (2018) and Coddington et al. (2016)) and the Reid and Schatten model (Christoudias (2016)) have significantly reduced long-term variability, as methodologically flawed reconstructions as equally probable alternatives to models, validated TSI models is scientifically unjustified. It creates a false equivalence that misrepresents the true state of solar irradiance research.

Quote given to Carbon Brief

Supporting evidence

1. [Lean, Judith L. Solar irradiance reconstruction. Living Rev. Sol. Phys. 15, 1 \(2018\). https://doi.org/10.1007/s41116-017-0009-6](#)
2. [Coddington, O., Lean, J. L., Pilewskie, P., Snow, M., & Lindholm, D. \(2016\). A solar irradiance climate data record. Bulletin of the American Meteorological Society, 97\(7\), 1265-1282.](#)
3. [Yeo, K. L., Krivova, N. A., Solanki, S. K., & Glassmeier, K. H. \(2014\). Reconstruction of total and spectral solar irradiance from 1974 to 2013 based on KPVT, SoHO/MDI, and SDO/HMI observations. Astronomy & Astrophysics, 570, A85.](#)
4. [Yeo, K. L., Solanki, S. K., Krivova, N. A., Rodriguez, J. C., Wenzler, T., Tapping, K. F., ... & Walton, D. \(2017\). Reconstruction of total and spectral solar irradiance from 1974 to 2009: Revised models and uncertainties. Journal of Geophysical Research: Space Physics, 122\(3\), 3888-3915.](#)
5. [Steinhilber, F., Beer, J., & Fröhlich, C. \(2009\). Total solar irradiance during the Holocene. Geophysical Research Letters, 36\(19\).](#)
6. [Egorova, T., Rozanov, E., Zubov, V., Karpechko, A., Manzini, E., Tschumi, J., ... & Peter, T. \(2018\). Chemistry-climate model SOCOL: version 3.0 with chemistry of VSLS and polar ozone depletion. Geoscientific Model Development, 11\(5\), 1863-1879.](#)
7. [Yeo, K. L., Ball, W. T., & Solanki, S. K. \(2020\). Are there multiple solutions for total solar irradiance and what do they mean for Earth's climate?. Space Science Reviews, 216\(8\), 1-19.](#)
8. [Connolly, R., Soon, W., Connolly, M., Baliunas, S., Berglund, J., Butler, C. J., ... & Zhu, G. \(2021\). How much has the Sun influenced northern hemisphere temperature trends? An ongoing debate. Research in Astronomy and Astrophysics, 21\(6\), 131.](#)
9. [Connolly, R., Soon, W., Connolly, M., Baliunas, S., Berglund, J., Butler, C. J., ... & Zhu, G. \(2023\). How much has the Sun influenced Northern Hemisphere temperature trends? An ongoing debate–Reply to comments. Research in Astronomy and Astrophysics, 23\(4\), 049501.](#)

3 Human influences on the climate

Page 11

MISLEADING

The IPCC has downplayed the role of the sun in climate change, but there are significant solar irradiance reconstructions that imply it contributed to recent warming.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research

This is a vague and potentially misleading statement. The IPCC AR6 adopted the solar irradiance forcing values recommended by Matthes et al. (2016). Of the CMIP6 models in both cases, the TSI and solar irradiance reconstructions selected were the most commonly used and recommended ones. The selection of the spectral TSI and SSI variation that these were 'dependent' while assigning a low TSI, without examining in further detail the impact that these 'dependencies' have on the climate response. Additionally, the choice of the solar irradiance reconstructions implying a low variability was not to be considered. The most scientifically robust irradiance reconstructions indicate that this contribution has been relatively small in the order of 0.1°C since 1850 and that the apparent impacts to implicitly reference the work of Connolly et al. (2021) and (2023), although only the former is cited in the report. Connolly et al. (2021) argued that a significant portion, or even most, of recent global warming could be attributed to solar irradiance changes, which is inconsistent with the vast majority of solar irradiance reconstructions, right with low long-term variability and right with high variability. To resolve this balance, they conducted a sensitivity analysis of models with strongly varying low-variability reconstructions.

This is not correct and was tested by Connolly et al. (2021):

- Hoyt and Schatten (1993) has been thoroughly discredited (see Christoudias (2016)). It includes arbitrary scaling parameters and even fabricated data, with some cycles seemingly copied from another one's data.
- Lean et al. (1995) is an early version of the NRL (Naval Research Laboratory) model that has since been superseded by more advanced models (Coddington et al. (2016), Yeo, Krivova, Coddington et al. (2014), and Yeo et al. (2017)).
- Reid et al. (2000) used a simplified linear conversion from cosmogenic isotope production (e.g., 10Be to TSI, without considering the nonlinear physics in the geomagnetic modulation, atmospheric transport processes, or climate-driven reservoir changes. Reid and Turney (2008) used a similar simplified approach. Both studies lack STOKE-TI (Spectral And Total Irradiance Reconstructions) (Yeo et al. (2014)), which show significantly lower long-term variability. Furthermore, Steinhilber and Beer (2013) revised the original Reid et al. (2000) reconstruction which was further updated in Steinhilber et al. (2009). Both versions were tested in Connolly et al. (2021) and are included in the ensemble.
- Shapiro et al. (2011), an older high-variability model, was effectively replaced by Egorova et al. (2018), which itself has been criticised by Yeo et al. (2020) for exaggerating long-term TSI changes. Egorova et al. (2018) was included five times in the Connolly et al. (2021) ensemble, including Shapiro et al. (2011).

Crucially, in all these high-variability models, the long-term TSI trends were not derived from observational constraints but were imposed as assumptions. Modern versions of the NRL model (Lean (2018) and Coddington et al. (2016)) and the Reid and Schatten model (Christoudias (2016)) have significantly reduced long-term variability, as methodologically flawed reconstructions as equally probable alternatives to models, validated TSI models is scientifically unjustified. It creates a false equivalence that misrepresents the current state of solar irradiance research.

Quote given to Carbon Brief

Supporting evidence

1. [Lean, Judith L. Solar irradiance reconstruction. Living Rev. Sol. Phys. 15, 1 \(2018\). https://doi.org/10.1007/s41116-017-0009-6](#)
2. [A Reassessment of Grandmunder Total Solar Irradiance Variations Since 1700. Solar Physics](#)
3. [Solar Irradiance Variability is Caused by the Activity Cycle and Not the Grand Cycle - AGU](#)
4. [Solar Irradiance Variability: Comparison of Models and Measurements - ADS](#)

3.1 Components of radiative forcing and their history

Page 12

MISLEADING

The IPCC underrates the change in the radiative forcing by the sun to be negligible, based on their preference for data reconstructions that imply minimal solar change since pre-industrial times.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research

The selection of TSI reconstructions for use in CMIPs and IPCC ARs was based on the scientific robustness and performance of the models, rather than the magnitude of TSI variability they imply (Matthes et al. (2017)). The two reconstructions adopted were SATIRE and NRLTSI (now referred to as NRLTSI2). These models were chosen because they represented the most advanced TSI reconstructions at the time (Matthes et al. (2017)). The most empirical, physics-based model, which NRLTSI is a proxy-based regression model, which have been extensively validated and demonstrated excellent agreements with direct satellite measurements of TSI, making them the most reliable choices for climate model studies.

Quote given to Carbon Brief

Supporting evidence

1. [CMIP6 – Data models – Solar Forcing for CMIP6 v3.2](#)

3.1 Components of radiative forcing and their history

Page 12

MISLEADING

A particularly thorny issue is the gap in TSI data between 1989 and 1996 due to delays in the launch of a satellite following the Space Shuttle Challenger disaster in January 1986.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research

It is an overstatement to characterise the "ACRIM gap" in pre-TSI measurements. Between the ACRIM-1 and ACRIM-2 satellite monitoring experiments there was a "particularly thorny" issue. While it does contribute to uncertainty in the long-term trends of TSI variability, it is not the only issue. Other issues include the drift of the HF3 TSI sensor during the ACRIM-1 period, resulting in a false trend (Willson & Mordvinov (2003), Christoudias et al. (2016) and Kopecky et al. (2008)).

Quote given to Carbon Brief

Supporting evidence

1. [A Reassessment of Grandmunder Total Solar Irradiance Variations Since 1700. Solar Physics](#)
2. [The Greatest Hoax of the Sun - Part - XXIII - Grandmunder Solar Minimum - Watts Up With That?](#)
3. [Solar Irradiance Variability is Caused by the Activity Cycle and Not the Grand Cycle - AGU](#)
4. [Solar Irradiance Variability: Comparison of Models and Measurements - ADS](#)

3.1 Components of radiative forcing and their history

Page 12

MISLEADING

Section 3.1.2

Dr Mads Schaldach, chief scientist, Science and Technology Corp.

The paper by Jenkins et al. (2021) did not include volcanic aerosol effects and thus underrated the strong sign of the long-term solar radiation forcing. Subsequent papers (including my own) agree that Hunga would produce slight cooling. The inclusion of the Hunga paper here suggests new uncertainty in the calculations of the models and data analysis paper that Hunga was incorrect (e.g. Schmidhuber et al. (2023) and Zhang et al. (2023)).

Quote given to Carbon Brief

Supporting evidence

1. [Modeling the Radiative Forcing and Atmospheric Temperature Perturbations Caused by the 2022 Hunga Tonga-Hunga Ha'apai Volcanic Eruption | Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)
2. [Hemispheric Asymmetry Partly Model Ensembles Simulations with Observations and Records Projections of Arctic Amplification, Temperature Differences and Ice Extents - ADS](#)

3.2 Future emission scenarios and the carbon cycle

Page 13

FALSE

Projections of future warming given significant uncertainties since the IPCC emissions projections have tended to overstate actual subsequent emissions. For the IPCC's RCP8.5 scenario, which assumed a very high rate of emissions growth, Hausfather and Peters (2020) found that, when corrected to per-capita values, the IPCC's projections overestimated near-actual emissions by around 100 percent (1%).

Dr Zeke Hausfather, research scientist, Berkeley Earth

While some past IPCC emissions scenarios have overstated future total CO2 emissions (RCP8, A1FI, RCP8.5), others have underestimated them (SRES B2, RCP6). In general, the scenarios have tended to bracket the range of actual emissions, with the 2000-2020 emissions near the high end of the RCP8.5 scenario and within the range in recent years. Notably both SP2 and RP3 scenarios were all variations of relatively high emissions scenarios that could include aggressive emissions mitigation.

Quote given to Carbon Brief

Supporting evidence

1. [Emissions are on course to forecast the worst-case scenario](#)
2. [CMIP5 - Global Carbon Budget 2020](#)

3.2 Future emission scenarios and the carbon cycle

Page 13

FALSE

Figure 3.2.1 from IAM report. CO2 emission trends of reference and concentration-stabilized scenarios. Source: Figure 1 in Hausfather and Peters (2020) and Figure 10.5 in Stocker et al. (2013).

Dr Zeke Hausfather, research scientist, Berkeley Earth

Figure 3.2.1 focuses primarily on emissions scenarios that predate the SRES scenarios. The SRES scenarios resulted in the figure on page 10 and 11 and, and those actually track CO2 concentrations quite well. The disagreement is primarily with 1970s and 1980s era models, however, as Figure 1.3 from Hausfather et al. (2019) notes the trend in total forcing in these early models is actually a bit too fast to their exclusion of non-CO2 GHGs.

Quote given to Carbon Brief

Supporting evidence

1. [Evaluating the Performance of Past Climate Model Projections - Hausfather - 2020 - Geophysical Research Letters - Wiley Online Library](#)

3.2 Future emission scenarios and the carbon cycle

Page 13

FALSE

The implausibility of the RCP8.5 scenario was examined by Burgess et al. (2021). The implausibility of RCP8.5 is explained by its temperature increase of 4°C by 2100 relative to pre-industrial, or a "worst case", but subject to genuinely implausible owing to the implausibility of the amount of fossil fuel that can be burned to cause it. RCP8.5 is the product of a model that projected strong demand trends in energy use and the most basic trends diverge shortly from those of the mainstream SSPs. Against this, all recent trends point to much lower projections of energy use for the coming decades. Pielke Jr. et al. (2022) further showed that the business as usual projected for RCP8.5 was not the bottom of the envelope of plausible BAU projections and the more recent Shared Socioeconomic Pathways (SSP) scenarios [more]

Dr. Zeke Hausfather, research scientist, Berkeley Earth:

The authors do not clearly differentiate between the implausibility of a particular emissions scenario like the high-emissions RCP8.5 and the likelihood that any of 4.5 watts per metre squared. The latter can come about even not just through emissions, but from climate sensitivity (see Hausfather and Peters (2020). RCP8.5 is quite unlikely in my view that we end up at 8.5W/m2 by 2100 even with large carbon cycle feedbacks, but it is entirely plausible to get to 8.5W/m2 emissions pathways. RCP8.5's high radiative forcing becomes increasingly plausible past 2100 if emissions continue.

Quote given in Carbon Brief

Supporting evidence

- Hausfather, Zeke; Carbon cycle feedbacks could make global warming worse - Carbon Brief
- Hausfather, Zeke and Peters, Glen; Emissions scenarios versus observational characterization of CMIP5 warming and climate
- Pielke Jr., Roger

Supporting evidence

- [2021 - Global Carbon Budget 2021](#)
- [ATL.BAS.ERF.2100.rcp45-85.html](#)

3.2 Future emission scenarios and the carbon cycle

Page 74 and 75

FALSE



Figure 3.12: Observed and projected CO2 emissions. Source: IPCC. IPCC scenarios and observations of annual CO2 emissions from fossil fuels and industry, 1990–2020. Note: the y-axis is in gigatonnes (billion tonnes) of CO2 per year.

Dr. Zeke Hausfather, research scientist, Berkeley Earth:

The authors mistakenly conflate total global observed CO2 emissions in this text with CO2 emissions from fossil fuels alone (which as shown in Figure 3.12). If we'd CO2 emissions from land-use change as well as fossil fuel emissions, then CO2 emissions are right in the middle of the SSP5-8.5 (on with SSP3-7.0 and SSP4-6.0, below SSP5-8.5 and SSP3-7.0, and above SSP2-4.5, SSP1-2.6 and SSP4-3.4.

Quote given in Carbon Brief

Supporting evidence

- [2021 - Global Carbon Budget 2021](#)
- [ATL.BAS.ERF.2100.rcp45-85.html](#)

3.2 Future emission scenarios and the carbon cycle

Page 75

MISLEADING

"Recent data from NOAA of carbon dioxide (CO2) in the Earth's atmosphere, almost all of it by the form of CO2 fuel- and land-use, dominated [plant growth and decay] and physical processes (ocean absorption and outgassing) have closely tracked the RCP8.5 trajectory through the Earth's surface through MOLE and the land and CMIP6, with the scenario. Below, however, activities become significantly different from the atmosphere-ocean roughly, between with emissions. Not learning fossil fuels (coal, oil, and gas) emissions reduced from the observed value is in the emissions range of the RCP8.5 scenario.

Dr. Daniel Cross, retired atmospheric physicist:

The total mass of carbon in the atmosphere is increasing over time as the carbon dioxide concentration increases. Crisp et al. (2022) quantified a mass of 877 Peta grams of Carbon (Pg C) over Pg C is equivalent to two billion tonnes of carbon (2,000 x 10^15 billion tonnes of CO2), which was a good estimate in 2020, when the concentration of CO2 dry air mole fraction (a measurement of concentration) was around 412ppm. That's around 412 molecules of CO2 per million molecules of dry air. The mass of the atmosphere is around 5 x 10^18 kg. The quoted ground-truth accuracy of carbon between land and the atmosphere and between the ocean and the atmosphere are estimates and their relative errors are reviewed. Here, they look at the exchanges of MOLE with the atmosphere and the oceans exchange 120GtC with the atmosphere. They quote the land exchange as 60GtC with the atmosphere (Ciais et al. (2022). That paper cites gross ocean-atmosphere carbon fluxes of 90GtC per year and gross land-atmosphere carbon fluxes of 120 GtC (GtC per year).

Then and this paragraph by noting that the 10.5GtC emitted by fossil fuel use and cement manufacturing is only about 5% of annual emissions by the land biosphere and the ocean. The 10.5GtC is 10.5 billion tonnes of carbon, or 38.5 billion tonnes of CO2.

Quote given in Carbon Brief

Supporting evidence

- [Note Well: On the Undetermined Net Land Ocean Atmosphere Carbon Cycle - Crisp - 2022 - Review of Geophysics - Wiley Online Library](#)

3.2 Future emission scenarios and the carbon cycle

Page 75

MISLEADING

The carbon cycle accommodates about 50% of humanity's annual anthropic emissions of carbon into the air by naturally incorporating it through plant growth and ocean absorption, while the remainder accumulates in the atmosphere (Ciais et al. (2013). For this reason, the annual increase in atmospheric CO2 concentration averages only about half of the [more]

Dr. Daniel Cross, retired atmospheric physicist:

Ciais, the report author recognise that existing observations show that natural processes in the land biosphere and ocean consistently remove about 50% of the anthropogenic CO2 emissions. This is the result of the net effect of many natural fluxes and fossil fuel emissions less forecast. They attribute the increasing land uptake to the "fertilization effect" of increased CO2 in the air (chapter 7.1 of the AR5 report). The chapter fails to recognise that the increasing leaf area index (LAI) does not always translate into increased plant growth. Crisp et al. (2022) show that in the tropical regions of the tropics, where the largest increases in LAI have been seen over the past 30 years, have started to demonstrate that CO2 sinks to an annual mean. Crisp et al. (2022) changes are due to the impacts of human activities (e.g. deforestation, forest degradation, land clearance, changes in farming practices) and natural effects (e.g. droughts).

The authors also fail to note that while tropical land is absorbing less CO2, these losses are being compensated mostly by increased carbon uptake by forests at mid and high latitudes. Chuine, Cheaib and Kunstler (2016) and Crisp et al. (2022) only examine one temperate woods and gets a result comparable to that found by Hasibeder et al. (2015), but does not do the depth. In the Hasibeder et al. (2015) study, it was found to see if the GSS has been effectively detected and removed (as Hasibeder et al. (2015) find that it was). The analysis in the Chuine, Cheaib and Kunstler (2016) assessments to ADS satellites global. Scanfield et al. (2019) and the US Climate Reference Network (US. Hasibeder et al. (2015) find that this the same sort of warming as the full land station network during the period of overlap, suggesting minimal UHI contamination. Chuine, Cheaib and Kunstler (2016) and Crisp et al. (2022) find that the net effect of the tropical forests is a net source of CO2 to the atmosphere. It is extended, would have a much smaller effect on resulting global surface temperatures.

Quote given in Carbon Brief

Supporting evidence

- [Quantifying the effect of afforestation on U.S. Historical Climatology Network temperature records - Hasibeder - 2015 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)
- [Evaluating the impact of U.S. Historical Climatology Network homogenization using the U.S. Climate Reference Network - Hasibeder - 2016 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)

3.3 Future emission scenarios and the carbon cycle

Page 75

MISLEADING

Section 3.3.2 substitutes six "CO2 uptake by land processes" and "CO2 uptake by ocean processes".

Dr. Daniel Cross, retired atmospheric physicist:

The last two subsections of this section focus on the carbon cycle models used by the Global Carbon Project. For both land and ocean, they note that this set of models predicts a range of estimates for the fraction of annual CO2 by the land and ocean carbon sinks. They use these models to first under the utility of these methods for projecting future CO2 concentrations but also as a means to calculate a budget. This is interesting because they fail to note that all of the Dynamic Global Vegetation Models (DGVMs) used in the analysis are from the same research group and that the authors of this document advocate as a key benefit of increasing atmospheric CO2 amounts.

Quote given in Carbon Brief

Supporting evidence

3.5 Simulating carbon–climate interactions

Page 78

MISLEADING

Paragraph 1 of the section 3.5 refers to trends between real and simulated atmospheric, although their definition of trends included local populations up to 10,000 persons with the relative influence of urbanisation (Hausfather and Peters (2020) and Hausfather et al. (2013).

Dr. Zeke Hausfather, research scientist, Berkeley Earth:

While Peterson et al. (1999) used a simple proxy for urbanity, the follow-up Hausfather et al. (2013) study used four definitions proxies for urbanity in the conterminous U.S. It found little residual urban heat island (UHI) bias in the homogenized NOAA data, even when only rural stations are used. Hausfather et al. (2013) also found that the process of warming in the full land station network during the period of overlap, suggesting minimal UHI contamination. Hausfather et al. (2013) find that the net effect of the tropical forests is a net source of CO2 to the atmosphere. It is extended, would have a much smaller effect on resulting global surface temperatures.

Quote given in Carbon Brief

Supporting evidence

- [Quantifying the effect of afforestation on U.S. Historical Climatology Network temperature records - Hasibeder - 2015 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)
- [Evaluating the impact of U.S. Historical Climatology Network homogenization using the U.S. Climate Reference Network - Hasibeder - 2016 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)

3.5 Simulating carbon–climate interactions

Page 78

MISLEADING

The influence of UHI warming is important in precipitation, in other words it is stronger in low population density than in high, and local urbanisation around the GHCN-CAMS [more]

Dr. Zeke Hausfather, research scientist, Berkeley Earth:

This is both overly simplified – the actual environment immediately around the station matters a lot more than regional population density – and incorrect as shown in the literature. Some studies using pretty strict cutoffs for urbanity (e.g. Parnikoz et al. (2015)) still find statistical difference in urban and rural stations.

Quote given in Carbon Brief

Supporting evidence

- [Quantifying the effect of afforestation on U.S. Historical Climatology Network temperature records - Hasibeder - 2015 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)

3.5 Simulating carbon–climate interactions

Page 78

FALSE

"In Germany, which there is clearly warming in the land record, there is also evidence that it is boosted upward by patterns of urbanisation and land-use change (Parnikoz et al. (2015)).

Dr. Zeke Hausfather, research scientist, Berkeley Earth:

The authors of the Dyck report make a strong claim based on papers by some of their own (McKitrick and Symon) and by other skeptic author studies that purport to show that there is a bias in the temperature record. But there is a meaningful controversial UHI bias in land temperature records. Hausfather et al. (2013) and Wickham et al. (2013) only examine raw temperature records and gets a result comparable to that found by Hasibeder et al. (2015), but does not do the depth. In the Hasibeder et al. (2015) study, it was found to see if the GSS has been effectively detected and removed (as Hasibeder et al. (2015) find that it was). The analysis in the Chuine, Cheaib and Kunstler (2016) assessments to ADS satellites global. Scanfield et al. (2019) and the US Climate Reference Network (US. Hasibeder et al. (2015) find that this the same sort of warming as the full land station network during the period of overlap, suggesting minimal UHI contamination. Chuine, Cheaib and Kunstler (2016) and Crisp et al. (2022) find that the net effect of the tropical forests is a net source of CO2 to the atmosphere. It is extended, would have a much smaller effect on resulting global surface temperatures.

Quote given in Carbon Brief

Supporting evidence

- [Quantifying the effect of afforestation on U.S. Historical Climatology Network temperature records - Hasibeder - 2015 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)
- [Evaluating the impact of U.S. Historical Climatology Network homogenization using the U.S. Climate Reference Network - Hasibeder - 2016 - Journal of Geophysical Research: Atmospheres - Wiley Online Library](#)

Part I: Direct impacts of CO2 on the environment

1.1 Components of radiative forcing and their history

Page 21

MISLEADING

Box 6.1 in the IPCC AR5 Report addresses the climate impact of volcanic eruptions, noting that explosive volcanic eruptions injected sulfur dioxide (SO2) into the stratosphere. This included the 1815 Tambora eruption that resulted in the "year without summer", and notable forced failures across the Northern Hemisphere. The report also addresses the role of the relatively small forcing due to the submarine volcano Hunga Tonga-Hunga Ha'apai eruption in January 2022 [more]

Prof Jonas Nycander, professor of climate science, Fusion National University:

This paragraph is misleading and does not accurately reflect assessment on the volcanic forcing in this AR6. For the models in WG1 Ch 7, as documented in WG1 Cross Chapter Box 6.1', Annex II, the model study made clear that this Cross Chapter Box to the WG1 Chapter 6 synthesizes assessments of volcanic aerosol forcing and their impacts on Earth system components across AR6 WG1 Chapters 2, 3, 4, 5, 6, 7, and 11. The summed text implies that the AR6 assessment of the volcanic forcing in the ensemble models orientation to relatively relatively small forcing. However, AR6+ particularly Section 2.2.5 provides a detailed assessment of volcanic aerosol forcing. Specifically, AR6 Chapter 2.2.2 notes: "a series of small to moderate eruptions since 2000 resulted in a negative forcing of about −0.1 W m−2 relative to 1850–1900 (Vernier et al., 2011; Andersson et al., 2015; Schmidt et al., 2018). To conclude, strong individual volcanic eruptions cause multi-annual reductions in radiative forcing, because of the large magnitude and variability of SAOD and its associated volcanic aerosol forcing since 1750 that are unusual in the context of at least the past 2.5 kyr (medium confidence).

Quote given in Carbon Brief

Supporting evidence

- [Chapter 2 - Climate Change 2021: The Physical Science Basis](#)

1.2 Future emission scenarios and the carbon cycle

Page 21

MISLEADING

Although the IPCC does not claim its emission scenarios are forecasts, they are often treated as such.

Prof Jonas Nycander, professor of climate science, Fusion National University:

The text seems to indicate that the IPCC is unclear about the meaning of emission scenarios, which is not the case. See Cross Chapter Box 6.1 in AR6 which explicitly states how emission scenarios are to be interpreted and used. For example, here is the description of the scenario used in this SPM: 1 of 3 AR6 synthesis report: "Modelled scenarios and pathways used to explore future emissions, climate change, related impacts and risks, and possible mitigation and adaptation strategies are based on a range of assumptions, including socio-economic variables and mitigation options. They are quantitative projections and not predictions and are neither forecasts nor policy neutral with regard to the assumptions underlying the scenarios in the literature assessed in this report, which do not cover all possible futures."

Quote given in Carbon Brief

Supporting evidence

- [SPM](#)

1.2 Future emission scenarios and the carbon cycle

Page 21

MISLEADING

The IPCC's development of new set of scenarios for AR6, the "Shared Socioeconomic Pathways" (SSPs), has been criticised for being too pessimistic in the RCP8.5 scenario [more]

Prof Jonas Nycander, professor of climate science, Fusion National University:

This text is false because the SSP scenarios were developed not by the IPCC, but by international science communities. It is important to recognized that the AR6 WG1 assessed a large number of plausible modelled scenarios, of which the SSPs were comprised based on their assessed global warming over the 21st century, and their representation of the full range of forcing in the literature. The SSP5-8.5 scenario was developed to explore the upper end of the range of forcing that exceed RCP8.5. The IPCC AR6 WG1 assessed the climate response to selected five scenarios from the SSPs, which were assessed to provide a representative sample from across the range of possible future development of anthropogenic drivers of climate change found in the literature. It is also very important to note that the IPCC AR6 WG1 did not comprehensively assessed feasibility of emissions scenarios including SSP1-1.9 and SSP1-1.9.

Quote given in Carbon Brief

Supporting evidence

- [SPM](#)
- [Chapter 7: Emissions: pathways, climate impacts, and economic](#)

Part II: Climate response to CO2 emissions

4.2 Model-based estimates of climate sensitivity

Page 57

FALSE

Section 4.2.1 does not refer to climate models based on the laws of physics, but to statistical models, relying instead on data-driven methods.

Anonymous,

Akike used multiple lines of evidence in the assessment of climate sensitivity: observational constraints on the climate system, paleoclimate evidence and model-based emergent constraints. Indeed, the authors of Sherwood et al. (2020) used their own climate-base paleographic term by saying: "For AR6, the IPCC placed primary weight on the results of Sherwood et al. (2020) who assessed multiple lines and paleoclimate/climate properties with the process-based approach." Sherwood et al (2020) used climate models as one of their lines of evidence.

Quote given in Carbon Brief

Supporting evidence

- [WG1 AR6 Working Group I: Support for Policymakers: Climate Change 2021: The Physical Science Basis](#)

4.3 Data-driven estimates of climate sensitivity

Page 58

FALSE

Most climate models simulate that rising GHGs will weaken the west-east temperature gradient, which led the IPCC to AR4 to conclude that data-driven ECS estimates underestimate the future ECS values. Recent studies (e.g. Gregory et al. (2019) pointed out that, in models, the west-east temperature gradient has been strengthening over time. [For example, Dessler et al. (2018) found that the strengthening of the gradient in models explains the characteristics of ocean heat uptake and that is the reason why the models simulate a weaker, a steeper temperature was recently made by Lee et al. (2021), who concluded that the increase in the strength of the observed trend reflects the response to increasing GHG loading in the atmosphere", in other words, GHG warming could lead to a future strengthening rather than a weakening of the temperature gradient. Increased efficiency of warming could, therefore, be underestimated by the observed ECS in a warming climate.

Dr Richard Tol, Emeritus Professor of Economics, Columbia University:

The report is right to point out that there is some attention to the fact that the observational warming record is not matching climate models relatively predict a weakening in response to rising GHGs. The report is also right that we suggested models get the response wrong due to biases in how they represent the tropics. However, the IPCC also underscores on the relation between EST, radiation and climate sensitivity for the short period of observations and the long-term response to the expected doubling of CO2.

Quote given in Carbon Brief

Supporting evidence

- [Lee et al. 2021 - Earth's Energy Imbalance Revisited](#)

4.3 Data-driven estimates of climate sensitivity

Page 58

FALSE

An argument emphasised in AR6 is that data-driven ECS estimates might underestimate the future warmingresponse to GHGs because of a so-called "pattern effect" (Forster et al. (2021). The Tropical Pacific Ocean is currently showing a La Niña-like pattern of warming, in which the Earth surface heat is spare, but atmosphere remove heat more effectively (Loeb et al. (2021). Climate models tend to simulate an El Niño response to GHG warming, which is why the IPCC assessing would underestimate short-term [more]

Prof Piers Forster, professor of climate science, University of Leeds:

The citation is in, but rather obtuse and abstruse, given the large amount of information that is contained in the section by the IPCC report - it's my chapter, I wrote it. I also find the literature in the section either what's the section, or you're missing a lot of other literature that is relevant to the section. I'm also finding it missing a lot of other literature with conflicting evidence. Lots of literature that is in the IPCC report and evidence published since – including my own work – suggests a significant and high climate sensitivity. If you are going to present a range of arguments to my IPCC chapter produced by many international authors and three rounds of peer review, including multiple expert reviewers, you would need to explain where these other studies we referenced in IPCC go wrong. I therefore find this argument weak.

Quote given to Gerbert Roe

Supporting evidence

- Chapter 7 The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity | Climate Change 2021: The Physical Science Basis

4.5 Data-driven estimates of climate sensitivity

Page 39

MISLEADING

A different argument was recently made by Jim et al. (2016), who concluded that, "the trajectory of the observed trend reflects the response to increasing GHG loading in the atmosphere". In other words, GHG warming should lead to a future strengthening (also) than a weakening of the temperature gradient. Increased efficiency of atmospheric cooling implies, if anything, that the future ECS for a warming climate might be lower than current estimates.

Prof Sukyoung Lee, *distinguished professor of meteorology*

The report cited my article out of context. Current climate models predict tropical Pacific sea surface temperature (SST) gradients that do not align with observational records, whereas the simple model used in my study does. The main reason that the climate models have no value for predicting the effects of human activities on future climate. My study simply identified a specific aspect where models may need to be improved to make the predictions more accurate.

Quote given to Gerbert Roe

Supporting evidence

- Quantifying the processes of accelerated convection Thomas Piatrou, recension: outside Section versus local Hadley Circulation Changes

5 Chapter summary

Page 51

FALSE

Climate models show warming times in many aspects of their reproduction of the last several decades. In response to increased greenhouse gas, this product has not been warming at the surface except in the models with lowest ECS.

Prof Stephanie Evring, *professor of climate modelling, German Aerospace Center & University of Bremen*

The CMIP6 historical simulations assessed in IPCC AR6 have an ensemble-mean global surface temperature change within the AR6 assessed very likely range for the historical period and observed warming is within the 5–95% range of the CMIP6 ensemble. Many of the models project the correct sign of the observed trend in tropospheric and lower-stratospheric temperature.

Quote given to Gerbert Roe

Supporting evidence

- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis

5.2 Surface warming

Page 53

MISLEADING



Figure 5.2: Global surface air temperature change (°C) relative to 1850–1900 simulated using only natural forcings (blue) and both natural and human forcings (red), and observed (black). Thick lines show the multi-model mean from CMIP6 and the thin lines show individual ensemble members. The CMIP6 multi-model mean is calculated from the same number of models as in Figure 5.1 and the observations are the same as in Figure 5.1. Anomalies are relative to 1850–1900 and are calculated from annual mean data. Sources: Ribes et al. (2021) for HadCRUT5 data; Haustein et al. (2017) for Cowtan & Way data; Morice et al. (2012) for GISTEMP data.

Figure 5.2 (Text only): Global surface air temperature change (°C) relative to 1850–1900 simulated using only natural forcings (blue) and both natural and human forcings (red), and observed (black). Thick lines show the multi-model mean from CMIP6 and the thin lines show individual ensemble members. The CMIP6 multi-model mean is calculated from the same number of models as in Figure 5.1 and the observations are the same as in Figure 5.1. Anomalies are relative to 1850–1900 and are calculated from annual mean data. Sources: Ribes et al. (2021) for HadCRUT5 data; Haustein et al. (2017) for Cowtan & Way data; Morice et al. (2012) for GISTEMP data.

Prof Stephanie Evring, *professor of climate modelling, German Aerospace Center & University of Bremen*

Figures 5.2 and 5.3 and related conclusions in the report need to be revised to use a higher reference period for the anomalies to ensure the influence of natural variability is minimised in model-observational comparisons (e.g. reference period 1850–1900 or 1861–1881). Also, the dates for the observed record to be used should be consistent (i.e. evaluated beyond the satellite era, and 23 accounts for observational uncertainty).

Quote given to Gerbert Roe

Supporting evidence

- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis

5.4 Vertical temperature profile mismatch

Page 57

MISLEADING

Figure 5.4: Vertical warming pattern for tropics (20S to 20N). Horizontal axis: Temperature. Source: Annotated version of IPCC AR5 Figure 9.SPM.6

Dr Katia Wenger, research scientist, department of climate modelling, University of Bremen & German Aerospace Center

The warming bias in the atmosphere's temperature profile is not "common", but can be found in certain periods and regions. Differences can, for example, be found in the 3rd simplified figure in IPCC AR5 SM 10.1. Models and observations were compared in Section 5.3.1.3 of the IPCC AR6 Report (Eyring et al. (2021)). Here, different periods are included, showing that the tropical atmospheric temperature trends from observational datasets and models agree well over the period 1979–2014. Only for about year 1998 to 2014, with observations showing weaker tropospheric warming compared to climate models.

Quote given to Gerbert Roe

Supporting evidence

- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis
- Chapter 2SM—Detection and Attribution of Climate Change: from Global to Regional - Supplementary Material | AR6

5.5 Stratospheric cooling

Page 58

MISLEADING

An important element of the expected general "fingerprint" of anthropogenic climate change is simultaneous warming of the troposphere and cooling of the stratosphere. This latter feature is also influenced by ozone depletion and recovery. ARs acknowledged that models had further observed that not until the year 2000. The stratosphere has shown some warming, since, scarcely to model projections.

Dr Jelle Kennedy, *scientist*

The poverty of their viewpoint is exemplified in the section on 'stratospheric cooling' about one-third of which consists of a long quote from the IPCC AR4 WG1 chapter 2, thought to cite a deceptively short paragraph from the IPCC AR5 WG1 Report, completely ignoring the section on the causes which can be found in Section 5.3.1.2.2 of the IPCC WG1 report. They can't even be bothered to finish the one additional quote that takes up much of the rest of the section, the one from Philipona et al. (2014). The quote, which is from the abstract but not that with a full stop, where the abstract goes on, after a comma, to say, 'which is consistent with a stronger from ozone depletion and greenhouse gas increases'. The full abstract is available for free, which is a nice touch and the report refers to the full paper in other parts of the report, but even the lowest little bit."

Quote from [redacted]

Supporting evidence

- Chapter 2: Changing State of the Climate System | Climate Change 2021: The Physical Science Basis
- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis
- Philipona, R., Dürr, B., Marty, C., Ohmura, A., & Wild, M. (2004). Fingerprint of the ozone hole: Warming, cooling, and a future recovery. Geophysical Research Letters, 31(16). https://doi.org/10.1029/2004GL020191 - Journal of Geophysical Research: Atmospheres - Wiley Online Library

5.4 Vertical temperature profile mismatch

Page 58

FALSE

The IPCC view did not assess this issue.

Dr Katia Wenger, research scientist, department of climate modelling, University of Bremen & German Aerospace Center

The warming bias in the atmosphere's temperature profile discussed in IPCC AR5 WG1 Chapter 5.3.1.2.1 (Eyring et al. 2021).

Quote given to Gerbert Roe

Supporting evidence

- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis

5.5 Stratospheric cooling

Page 58

FALSE

A combination of tropospheric warming and stratospheric cooling is a commonly cited "fingerprint" of anthropogenic climate change. Stratospheric warming since 2000 is consistent with continued ozone and greenhouse warming, a prediction that is not found in climate model simulations and is not supported consistent with the anthropogenic fingerprint.

Dr Benjamin Santer, *atmospheric scientist, Lawrence Livermore National Laboratory*

The DoE claim is not true. Climate models can and do show recovery of lower stratospheric temperature after 2000 in response to ozone recovery (and to accord with satellite observations). Over the full satellite era (1986 to 2016), models show very large ranges of the amplitude of upper-stratospheric response to human-caused changes in CO2 and ozone (see Fig. 1 in the appended paper). The observed vertical structure of tropospheric warming and stratospheric cooling is also in good agreement with model results. The full report cites the appended paper (Santer et al. 2013) and flags up as evidence of absence" of a human fingerprint on the vertical structure of atmospheric temperature. This 2013 paper actually provides unique evidence FOR the positive identification of the human fingerprint in satellite data.

Quote given to Gerbert Roe

Supporting evidence

- Exceptional stratospheric contribution to human fingerprints on atmospheric temperature - PNAS

5.8 US corn belt

Page 61

FALSE

One of the largest discrepancies between models and observations is in the US corn belt, for which 47 particular importance to global food production. Figure 9.9 shows the various trends for summertime (June, July, August) for the 12 states east-west (IN, IA, IL, MO, OH, WI, MI, MN, SD, ND) during 1979–2012. Only 10 of the ensemble trends reach the observed trend.

Dr Christine Delsole, *lead author, IPCC AR6 WG1*

There is no source for this and no methodological description. I don't believe the comparison between models and observation is correct.

Quote given to Gerbert Roe

Supporting evidence

- Chapter 3: Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis

5.9 US corn belt

Page 61

FALSE

An illustrated in Chapter 9, the anticipated negative effects of atmospheric temperature at 10 corn yields have not materialized, in contrast to widely published studies proclaiming that detrimental future impacts are already being experienced (e.g., Burgess et al. (2016)).

Dr Richard Seager, Ewing Lamont research professor, Columbia University

Our 2018 paper referred to changes in utility and implications for what crops are grown and farm size and not what crops yields. We suggest that changes patterns and values of precipitation and potential evapotranspiration would increase utility over most of the US, which would, in the absence of adaptation, restrict cultivation to regions further east. But neither part of the two part paper claims yields are reducing due to climate change.

Quote given to Gerbert Roe

Supporting evidence

- Whither the 100th meridian? The once and future physical and human geography of America's arid-humid divide. Part II: The multidecadal past - Earth Interactions

6 Extreme weather

Page 65

Section 6

Prof Kerry Emanuel, *professor emeritus of atmospheric science, Massachusetts Institute of Technology*

The authors begin sensibly, in the introduction, stating that: "Climate is always the statistical properties of weather over decades, not single events. Further, there are only about 150 years of reliable observational data on which to assess changes in climate." But belief interval does not begin to contain all the extreme events that the climate system can create on this time. It is odd, then, that they proceed to present descriptions of extra shorter length and claim not to find evidence of extreme events in them, thereby leading the reader to conclude that there are no such extreme events. The authors should start with their original statement and made the correct conclusion that observations usually do not suffice to detect extreme events because of the predicted magnitude, and then turned to an exposition of what theory and models have to say.

It is very odd that there is almost no mention of theory and models [in] extreme events. The authors should explain that the reason for the lack of observed trends is virtually all of the tropical cyclone proxies register storms. And, contrary to the statement in the Draft report, an upward trend in the proportion of very strong hurricanes/tech been observed and published. But instead of using any of this, these authors relate their own introductory statement by citing short and generally unreliable records. For example, they claim that there seems to be a detectable trend in continental US hurricane landfalls. But at an average of three landfalling storms per year, there are not enough storms to detect a trend. The most reliable measurements of landfalling storms in Caribbean region had a high population density (and associated newspaper accounts) going back to the early 19th century. They could have looked at, but didn't, and they did not look the US. Had there done so, they would have discovered a clear upward trend.

Quote from [redacted]

Supporting evidence

- Chapter 11: Weather and Climate Extreme Events in a Changing Climate | Climate Change 2021: The Physical Science Basis

6 Extreme weather

Page 65

Section 6

Prof Sonia Seneviratne, *head of Institute for Atmospheric and Climate Science, ETH Zurich*

The text of chapter 6 cites often chapter 11 of the IPCC AR6, but is a clear example of cherry-picking. The authors repeatedly highlight low-confidence statements from the IPCC AR6 chapter 11 on changes in data extremes which are mostly on side topics of little relevance, but they omit the very high-confidence statements from chapter 11.

Quote given to Gerbert Roe

Supporting evidence

- Chapter 11: Weather and Climate Extreme Events in a Changing Climate | Climate Change 2021: The Physical Science Basis

6 Extreme weather

Page 65

Section 6

Most types of extreme weather exhibit no statistically significant long-term trends over the available historical record.

Prof Sonia Seneviratne, *head of Institute for Atmospheric and Climate Science, ETH Zurich*

Importantly, there is no evidence substantiating the claim of the authors of this SOD. Indeed, Chapter 11 of the IPCC AR6 WG1 report assessed that there are significant long-term trends over the available historical record". It is not clear which types of 'extreme weather' the authors refer to, which requires they are referring to, and what is the demonstration underlying the claim that these trends are not significant or statistically significant long-term trends. I would ask the authors to provide the arguments substantiating this claim, as there is no rationale provided for this statement in the chapter.

I would also highlight the following conclusions from the IPCC AR6 chapter 11 which are relevant to the statement in question:

- "It is re-established fact that human-induced greenhouse gas emissions have led to an increased frequency and/or intensity of some weather and climate extremes since pre-industrial time, in particular for temperature extremes."

- "Human influence has likely contributed to the observed global scale changes in the frequency and intensity of daily temperature extremes since the mid-20th century."

- "The frequency and intensity of heavy precipitation events have likely increased at the global scale over a majority of land regions with good observational coverage. Heavy precipitation has likely increased on the continental scale over three continents (North America, Europe, and Asia."

- "More regions are affected by increases in agricultural and ecological droughts with increasing global warming (high confidence)."

- "It is likely that the global proportion of Category 3–5 tropical cyclone instances has increased over the past few decades."

Quote given to Gerbert Roe

Supporting evidence

- Chapter 11: Weather and Climate Extreme Events in a Changing Climate | Climate Change 2021: The Physical Science Basis

6.2 Temperature extremes

Page 67

Section 6.2

Prof Erich Fischer, lecturer at the department of environmental systems science, ETH Zurich.

The section on hot extremes is an extreme example of cherry-picking results. The report fails to acknowledge that the highlighted regions are some of the very few regions globally where the annual maximum temperature has not increased (see e.g. IPCC AR6 WG1 Fig. SPM1). The report further highlights that in some regions the number of hot days has not increased since the 1950s, but fails to mention that increases increased over the same regions and periods (see figure 2.1 in the fifth IUTA National Climate Assessment). Furthermore, the interpretation of the narrowest nights shows a positive trend over records decades (Singh et al. 2021). The report further fails to refer to the extensive literature discussing the role of land use changes and irrigation (Mueller et al. (2016), around Kerning (Mueller et al. (2017) and enforced internal variability (Singh et al.) in the regions with little trends.

Quote given to Gerbert Roe

Supporting evidence

- Warm hills Will | Climate Change 2021: The Physical Science Basis
- ERR
- Correlation decomposition extremes intensification over the Midwest USA and amplified over Western Europe | Communications Earth & Environment
- Causes of US Midwest summer temperature and precipitation extremes recently | Nature Climate Change
- Timing and magnitude of the United States' warming hole – IOPscience

6.8 Wildfires

Page 76

MISLEADING

Wildfire have been increasing on the years 1982 interval the incidence of this for the US is not increasing. The areas burned did increase but only until about 2007.

Prof A Park Williams, hydroclimatologist, University of California.

The report's section on wildfire lumps the entire US together, smearing out what's going on in the western ecoregions that dominate the fact that it is the rapidly increasing western US wildfire activity that accommodates this report's wildfire section in the first place. The report's statement that "the area burned by wildfires in forests of the western US has increased since the 1970s" is misleading. It is true that western forests have consistently shown wildfire trends in the US. In the western US, the annual area burned has tripled since the 1970s, and the area burned in 13 mid-elevation forests in the western US has quadrupled.

FALSE

Evaluating AR6 projections to 2020 with reference to the baseline period (1995–2014) shows that while most but not all models (~80%) will run hotter, a range of models will run cooler.

Prof Stefan Rahmstorf, professor of physics at the oceans, University of Potsdam

Stefan Rahmstorf's analysis, as mentioned above, finds that assuming that in the first half of the period half of the projected rise should have occurred. But, of course, the IPCC projections account for acceleration, so the projected rise is not linear but accelerated and rises. In fact, projecting sea level rise until 2050 can be extrapolated by the observed rate, including the observed acceleration, and projected rise to 2050 from the projections even though these are made in a completely different way.

Looking at the data, for the intermediate term (2015, the AR6 predicts a best estimate 9–10cm, relative to the same base period (1995–2014) (Table 9.9 of AR6 WG1). The AR6 predicts a rise of 15–20cm for the intermediate term (2030, the 10–year mean, that will be 10cm by 2050, well within the 15–20cm projected by the AR6.

Quote given in Question 4(b)

Supporting evidence

- [Sea Level Rise Projections Supported by Extrapolation of Tide Gauge Observations - Nunn, P.D. and Hosking, Anthony Glenn - Wiley Online Library](#)
- [Wikipedia](#)
- [wikipedia - Sea Level Research Group](#)

7 Changes in sea level

Page 71

FALSE

A single group models

Prof Stefan Rahmstorf, professor of physics at the oceans, University of Potsdam

Most US tide gauges show acceleration, so the east coast that is statistically significant and larger than the global average acceleration, so the west coast too so.

Quote given in Question 4(b)

Supporting evidence

- [Sea Level Rise in the U.S.: Open Mind](#)

7.1 Global sea level rise

Page 71

FALSE

Following the end of the Little Ice Age in the mid-19th century, tide gauges show that the global mean sea level began rising during the period 1830–1860, well before most anthropogenic greenhouse gases were emitted.

Prof Stefan Rahmstorf, professor of physics at the oceans, University of Potsdam

While it is true that that before 'most' of greenhouse gas emissions, it is often to compare the "start" of the rise with "most" of emissions. The industrial era is often assumed to start at ~1750 and CO2 concentration also starts to rise noticeably around 1830–1860, around the same time as sea level rise the briefing notes).

Quote given in Question 4(b)

Supporting evidence

- [Timeline of atmospheric carbon dioxide - Wikipedia](#)
- [Industrial Era - Wikipedia](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Models infer that natural internal drivers since 1850–1900 have changed global surface temperature by -0.1C to +0.1C, and internal variability has changed it by -0.2C to +0.2C, so the average having essentially no net effect on the observed warming.

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

As AR6 report presents this in Fig. SPM.2 as "Aggregated contributions to 2010–2019 warming relative to 1850–1900". This refers to the net effect of various climate drivers on global surface temperature between these two periods, not the maximum changes observed. The accompanying range represents the uncertainty in these estimates. It is nonetheless safe to assume that the net effect of natural internal drivers was small between 1850–1900, because the term is not given further scrutiny.

Quote given in Question 4(b)

Supporting evidence

- [wikipedia - Sea Level Research Group](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Based on Hawkins, this internal contribution of natural variability has been disputed by several publications that question the magnitudes of solar variability and internal variability from large–scale ocean circulations.

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

This claim is supported by only a small number of studies that rely on theory–driven, rather than empirically–replicable TSI reconstructions (e.g., Connolly et al. (2021) and (2023), Scafetta (2021a, 2021b, 2021c), Gao et al. (2021), and Lean and Rind (2023)). All of these studies rely heavily on the Hoyt and Schatten (1993) TSI series, which has been extensively criticised for its reliance on flawed sunspot data (due to arbitrary adjustments, methodological flaws and use of fabricated data. As such, this claim is not reflect a broadly accepted position in the field. The claim that Connolly et al. (2023) provides a comprehensive and up–to–date overview of our current understanding of long–term solar irradiance variations and also misleading as the practices underlying such claims are scientifically flawed. I note that although the Connolly et al. (2023) paper was published in 2023, the data it relies on terminated at the southward and irradiance season and its core conclusion is dated.

Quote given in Question 4(b)

Supporting evidence

- [WG1 AR6 Working Group 1 Summary for Policymakers | Climate Change 2021: The Physical Science Basis](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Models infer that natural internal drivers since 1850–1900 have changed global surface temperature by -0.1C to +0.1C, and internal variability has changed it by -0.2C to +0.2C, so the average having essentially no net effect on the observed warming.

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

As AR6 report presents this in Fig. SPM.2 as "Aggregated contributions to 2010–2019 warming relative to 1850–1900". This refers to the net effect of various climate drivers on global surface temperature between these two periods, not the maximum changes observed. The accompanying range represents the uncertainty in these estimates. It is nonetheless safe to assume that the net effect of natural internal drivers was small between 1850–1900, because the term is not given further scrutiny.

Quote given in Question 4(b)

Supporting evidence

- [Lean, Judith Hoyt, Douglas Rind, David Warm Years Enhanced by the Combined Effects of Anthropogenic Greenhouse Gases and Natural Variability](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Natural variability, the internal contribution of natural variability has been disputed by several publications that question the magnitudes of solar variability and internal variability from large–scale ocean circulations.

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

This claim is supported by only a small number of studies that rely on theory–driven, rather than empirically–replicable TSI reconstructions (e.g., Connolly et al. (2021) and (2023), Scafetta (2021a, 2021b, 2021c), Gao et al. (2021), and Lean and Rind (2023)). All of these studies rely heavily on the Hoyt and Schatten (1993) TSI series, which has been extensively criticised for its reliance on flawed sunspot data (due to arbitrary adjustments, methodological flaws and use of fabricated data. As such, this claim is not reflect a broadly accepted position in the field. The claim that Connolly et al. (2023) provides a comprehensive and up–to–date overview of our current understanding of long–term solar irradiance variations and also misleading as the practices underlying such claims are scientifically flawed. I note that although the Connolly et al. (2023) paper was published in 2023, the data it relies on terminated at the southward and irradiance season and its core conclusion is dated.

Quote given in Question 4(b)

Supporting evidence

- [Lean, Judith Hoyt, Douglas Rind, David Warm Years Enhanced by the Combined Effects of Anthropogenic Greenhouse Gases and Natural Variability](#)

8.3 Attribution of global warming

Page 69

MISLEADING

AR5 concluded that the best estimate of radiative forcing due to Total Solar Irradiance (TSI) changes over the period 1750–2011 was very small (0.05 W/m2, Myhre et al. 2013).

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

This is an example of imprecise wording that could be misleading. In AR5, the solar radiative forcing difference between the CMIP and RRR activity minima was estimated using TSI reconstructions from Wang et al. (2005), Steinhilber et al. (2009) and Krivova et al. (2010). These studies estimate a range of 0.05 to 0.071 Watts per square metre (W/m2) for the difference (based on seven year running means, with the extent that the AR5 report refers to Steinhilber et al. (2009), which has a five–year resolution and uses 1965 as the minimum). Based on these estimates, AR5 adopted a range of 0.1 W/m2 (uncertainty of 0.05 W/m2) for the change in solar radiative forcing since 1750. In the report, however, this defined solar forcing in a different way to reflect the difference between the average TSI for 1745–1765 and the average TSI for 1995–2014 (Myhre et al. 2013). This change in methodology is not acknowledged in the full report, which may lead to misunderstandings or incorrect comparisons between AR5 and AR6.

Quote given in Question 4(b)

Supporting evidence

- [IPCC](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Some acknowledgers substantially higher values and a much larger range of estimates of changes in TSI over the last several centuries, stating that the TSI between the Maunder Minimum (1645–1715) and the second half of the 20th century increased by 1.3 W/m2, a range that includes both low and high variability TSI data sets (Judith (2012)).

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

The sentence is vague in particular as to what they refer with 'higher values') and potentially conflates two related but distinct concepts: the magnitude of TSI differences, which were both mentioned but not clearly distinguished. Considering that the previous sentence discusses relative TSI variations, one can assume that the 'higher values' they refer to radiative forcing too, which would make the statement false. This is easily understood by comparing the differences in AR6 and AR5 about the range of solar radiative forcing:

- The AR5 WG1 report says TSI–source:
"In contrast to AR5, the solar effective radiative forcing (ERF) in this assessment uses the full solar cycle rather than solar minima. It has increased by 0.05 W/m2 since the start from all complete solar cycles from the start of the 1745 (CE1745–8) to recent (CE1995–2008) (CE1745–1765). The mean TSI data from 1995–2014 is 0.9 W/m2 higher than the assessment period for 2019. The best estimate solar ERF is assessed to be 0.01 W/m2, using the 14C reconstructions from SATIRE–M, with a likely range of 0.00 to +0.08 W/m2 (medium confidence). The uncertainty range is adopted from the evaluation of the maximum between the second half of the 20th century and the average TSI over the Maunder Minimum in 2019 using several datasets, resulting in an ERF range of 0.1 W/m2 (Figure 8.13). Solar forcing is assessed to have increased over the period over the period of reduced solar activity in the 17th century relative to the period 1750 to 2019 (medium confidence)." The uncertainty in the AR5 report (page 689) states:
"The best estimate from our assessment of the 14C data is a change in solar forcing between seven–year running mean radiative forcing (RF) between the minima of 1745 and 2008 of 0.05 W/m2. The assessment of the range of RF from TSI changes is 0.0 to 0.10 W/m2."

Thus, the solar radiative forcing estimates in AR5 and AR6 are broadly consistent, with only best estimate slightly lower. If the discussion is about TSI variations, then AR6, this would also be misleading. The AR6 WG1 page 397, box 7.01 differences between the Maunder Minimum and the second half of the 20th century is 1.3 W/m2. The lower bound likely corresponds to the SATIRE–T reconstruction, while the upper bound to SATIRE–M. Both of these are theory–driven reconstructions and both of the ones. It important to note that the TSI variations between the LIA and the Maunder Minimum but rather provides a theoretical constraint on the estimates. Empirical reconstructions, for instance, typically exclude TSI reconstructions with large amplitude variability, such as Figures et al. (2010). Almost all current TSI reconstructions indicate a significant change only from 1915 (Usoskin and Kovaltsov et al. (2021)). These include the SATIRE and NRL (now NRL) models reconstructed for CMIP6 and AR6.

Quote given in Question 4(b)

Supporting evidence

- [Lean, Judith Hoyt, Douglas Rind, David Warm Years Enhanced by the Combined Effects of Anthropogenic Greenhouse Gases and Natural Variability](#)
- [Chapter 7 | The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity | Climate Change 2021: The Physical Science Basis](#)

8.3 Attribution of global warming

Page 69

FALSE

The IPCC has only minimally discussed solar influences on global and regional climate.

Prof Mike Lockwood, professor of space environment physics and president of the Royal Astronomical Society, University of Reading.

Contrary to this statement, there is much discussion in the last three IPCC assessment reports.

Quote given in Question 4(b)

Supporting evidence

- [Chapter 7 | The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity | Climate Change 2021: The Physical Science Basis](#)

8.3 Attribution of global warming

Page 69

MISLEADING

Section 8.3.3

Prof Andrew Schreiber, college of Earth, ocean and atmospheric science Prof Veronika Eyring, professor of climate modelling and Prof Gerd Steinle-Neumann, professor of theoretical physics

Overall, this section overemphasizes uncertainty and claims a misleading overall lack of natural climate variability in IPCC AR6, whereas we do know with high certainty that anthropogenic greenhouse gas emissions have caused the global warming over the last 50–100 years. There are no credibly findings other than greenhouse gases that could have caused that warming. Temperature measurements in the ocean show that the ocean is warming from the top down, which is inconsistent with a role of ocean circulation. Solar irradiance has not increased since the 1950s, and the sun is in a slightly cooling conservation regime that if some irradiation had caused surface warming there must be cooling in the stratosphere. However, this is not observed. The ocean has warmed at all depths, most at the surface, and the stratosphere has cooled. The anthropogenic warming. This should be mentioned in the part entitled "Natural Variability of Large Scale Ocean Circulation" in the expert is misunderstood. Section 8.3.3. Although state of the art climate and Earth system models tend to underestimate natural internal climate variability on decadal timescales, climate models nonetheless have provided a pathway to better quantify its range. Using several such models, Hawkins et al. (2020) showed that the 15–year trend in global mean surface temperature (GMST) could reflect an interpretation or approximately +0.2 °C (90%CI: 0.0 to +0.4 °C) warming from about +0.2 °C (CanESM2) to +0.5 °C (EC-Earth3). This model-based estimate is consistent with observation-based estimates. For example, the following changes must be made in the IOM CMIP5 document: ". Revise the statements that claim an inadequate assessment of internal climate variability by IPCC AR6. IPCC AR6 has extensive peer–reviewed literature that led to the following assessment: "When estimated from observations, the observed 15–year trend in GMST is consistent with variability in global surface warming of +0.25°C to +0.25°C is small compared to the warming of about 1 °C observed during this period, which has been primarily attributed to the human influence." Include evidence from ocean warming patterns, which is inconsistent with changes in ocean circulation causing the observed global warming, but which is consistent with anthropogenic warming causing heat uptake from the top down. Include evidence from the lack of warming in the stratosphere, which is inconsistent with changes in the sun's irradiance causing the observed warming. Include evidence from the lack of increase in solar irradiance over the historical period, which has important implications for attributing recent trends to anthropogenic warming.

Quote given in Question 4(b)

Supporting evidence

- [Projections of Human-Induced Warming from the World's Oceans | Science](#)
- [Recent Surface Ocean Warming Contrasts Early 20th Century Attributable to Anthropogenic Forcing | Geophysical Research Letters - Wiley Online Library](#)
- [Chapter 9 | Human Influence on the Climate System | Climate Change 2021: The Physical Science Basis](#)

8.3 Attribution of global warming

Page 69

MISLEADING

There is substantial evidence for high solar activity in the second half of the 20th century (going to 1995) and extending into the 21st century, but a decline in the early 21st century; this period is often termed the "Modern Maximum" (Solanki et al. (2023)). Solanki (2013) and Lockwood et al. (2014). However, some scientists have concluded that it is not possible to be confident of any multi–decadal trend in TSI (Schrijver (2015)).

Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

Although the sentence in which this text is in factually accurate in isolation, the surrounding text creates a misleading impression. More importantly, our review paper, which they cited, explains why that subsequent paragraph has misleading and wrong statements."

Quote given in Question 4(b)

Supporting evidence

- [Lean, Judith Hoyt, Douglas Rind, David Warm Years Enhanced by the Combined Effects of Anthropogenic Greenhouse Gases and Natural Variability](#)

8.3 Attribution of global warming

Page 69

MISLEADING

However, the recommended forcing dataset for the CMIP6 climate model simulations used in the AR6 synthesis underestimates solar variations and has now been superseded (Matthes et al. (2017)).

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

- [SORCE - Total Irradiance - Daily Total Value for CMIPV6 v3.0](#)
- [The impact of solar irradiance variations | Nature Astronomy](#)

8.3 Attribution of global warming

Page 69

MISLEADING

While AR6 shows a substantially greater solar impact than does AR5, the overall impact of solar forcing on the climate was still assessed to be small compared to anthropogenic forcing.

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

The statement is vague and unclear in what it means by "AR6 shows a substantially greater solar impact than AR5", especially as the previous sentence refers to the fact that solar forcing was assessed to be small compared to anthropogenic forcing, implying a correspondingly small solar impact. The statement is misleading in that it implies confusion on the part of the AR6 authors regarding the distinction between radiative forcing and total solar irradiance (TSI) difference as presented in AR5 and AR6, as discussed in previous comments.

Quote given in Question 4(b)

Supporting evidence

8.3 Attribution of global warming

Page 69

MISLEADING

However, the impact of solar variations on the climate is uncertain and subject to substantial debate (Lockwood (2012); Connolly et al. (2021)) — something that is not evident in the IPCC assessments (Schrijver (2015)).

Dr Theodoros Chatzinikolaou, astrophysicist, New Plans Institute for Solar System Research.

Alls acknowledges the high variability reconstruction by Egerova et al. (2018) and references studies by Yeo et al. (2020) and Lockwood & Ball (2020), discussing the difficulties in reconciling solar variability with observed climate variations. However, the earlier study by Lockwood (2012) is not included because its findings are superseded by the more recent Lockwood et al. (2014) and Lockwood et al. (2016) studies, which are not listed here or AR6. This IPCC AR6 did not discuss the solar forcing uncertainty in irradiance modelling and, thus, its effects on Earth's climate. However, what the IPCC AR6 does not discuss is the fact that the solar forcing data set used in AR6 (from the Connolly et al. (2021) study). It is worth reporting that the Connolly et al. (2021) study is not a TSI reconstruction but rather a model of solar irradiance. Connolly et al. (2021) clearly pointed to TSI reconstructions, which are not empirical.

8.3 Attribution of global warming

Page 33

MISLEADING

There are several real composite TSI datasets that disagree as to whether TSI increased or decreased during the period 1986–96.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

It is also important to note that the effect of ACRIM gap in the trends of the different TSI composites is generally small (Chatzistergos et al. 2020, Yeo et al. 2020). While the issue with ACRIM data contributes to the uncertainty in the long-term trend of direct TSI measurements, a growing body of evidence points to the lack of a TSI increase during the ACRIM gap, evidence which the IAR report entirely overlooks. Notable examples include Scafetta & Willson (2013), Chatzistergos et al. (2021) and Krivova et al. (2009). So having given such emphasis on the ACRIM gap the IAR report gives the misleading impression that there is only one way to interpret TSI data, which could account for the global warming over the same period. Thus, this statement can be misleading.

Quote given to Carbon Brief

Supporting evidence

- Long-term changes in solar activity and irradiance - ScienceDirect
- Reconstructions of the Sunspot Number: Status Report - Solar Physics
- The Planetary Scale of the Sun – Part – XXII – Geophysical Research Letters – Wiley Online Library

8.3 Attribution of global warming

Page 33

MISLEADING

Neither, the satellite record of TSI is used to calibrate proxy models that infer past solar variations from sunspots and cosmogenic isotope measurements (Krivova Reference et al. [2010]).

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

There are several issues with this sentence, particularly concerning the choice of the sole reference, which only mislead readers into believing that existing total solar irradiance (TSI) reconstructions depend on the satellite record for calibrating and determining long-term trends. This is generally incorrect. Moreover, the phrasing gives the impression that only one reference exists, overlooking the important class of physics-based semi-empirical models.

1) Missing relevant literature: The statement omits key papers on irradiance reconstruction models (e.g., Coddington et al. 2016; Wu et al. (2018), and Chatzistergos et al. (2022a) and authoritative reviews (e.g., Chatzistergos et al. (2021) and 2024), and builds a skewed picture by citing only a more accurate and comprehensive perspective.

2) Choice of reference: The IAR cites only Velasco Herrera et al. (2015) to support the claim, which is problematic. That work is not an irradiance reconstruction model but a machine learning extrapolation dependent on the SATIRE-S model for its training series. Given the chaotic nature of solar activity, such extrapolations from limited data can naturally reach non-physical values. This is explicitly noted by Velasco Herrera et al. (2015) for 2011–20 as already contradicted by actual direct TSI measurements obtained since 2003.

3) Proxy models: These reconstruct irradiance variations by scaling or regressing solar activity indices against reference TSI data. While the choice of reference series can influence the magnitude of long-term changes in the reconstructed, the effect is not always big. For example, Chatzistergos et al. (2020) demonstrated that the choice of TSI reference has a relatively small impact on reconstructions of the solar contribution to global warming (see Scafetta, Figure 5 in Chatzistergos et al. (2021a) shows only a minor effect of TSI references on their relative contributions).

4) We have not only proxy models but also physics-based semi-empirical models, such as SATIRE-S and SSPM, which are entirely independent of satellite TSI composites to direct TSI measurements. However, this testing does not significantly affect the long-term trends, as shown in multiple studies (e.g., Chatzistergos et al. (2021), Yeroshi, Chatzistergos et al. (2023) explicitly states about their irradiance reconstruction that "*the TSI record is not used in the updated SATIRE-S TSI composite is independent of measured TSI*".

Quote given to Carbon Brief

Supporting evidence

- Understanding the secular variability of solar irradiance: the potential of CLA-S observations | Journal of Space Weather and Space Climate
- Reviews of the SATIRE-S irradiance reconstruction model of TSI: Effects of instrumental and SATIRE-S model changes | Astronomy & Astrophysics (A&A)
- Modelling solar irradiance from sunspot band photometric observations | Journal of Space Weather and Space Climate
- Reconstructing solar irradiance from historical Ca II K observations – A Method and its validation | Astronomy & Astrophysics (A&A)

8.3 Attribution of global warming

Page 33

MISLEADING

There is substantial evidence for high solar activity in the second half of the 20th century according to 1990's and continuing today (e.g. 1996, Solanki & Krivova) that it was higher than any period in either around the "Modern Maximum." (Chatzistergos et al. (2020), Scafetta and Willson (Scafetta et al. 2007).

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

The citation of Chatzistergos et al. (2021) in support of the statement about the "Grand Modern Maximum" of solar activity is somewhat misleading. First, the cited paper is a review-based exposition on solar irradiance modelling and long-term trends in irradiance reconstructions. The Grand Modern Maximum is mentioned only briefly in the context of sunspot number records (Section 2.1.1), and the paper does not assess or quantify the period's solar activity. The statement is not direct research output from the paper, discussed in results such as Chatzistergos et al. (2017) and Usoskin et al. (2016), which clearly stated its characteristics at all 27 solar cycles from 1700 to 2015. Other key studies cited in the IAR, Solanki et al. (2004) and Usoskin et al. (2007), followed this same approach. Second, the statement misrepresents the cited paper's conclusions. Whereas these data do not cause the full extent of the modern maximum, and the connection between irradiance and sunspot number records is not straightforward. Third, reconstructions like Wu et al. (2018), where both Sami K. Solanki and Ilya G. Usoskin are co-authors continue to support the existence of a Grand Modern Maximum. This suggest it was less pronounced than earlier records.

To present the authors in such a way that the statement was used to reconstruct irradiance variations and returned a rather small secular trend, which would conflict with the claim they have made. This is important to recognise that the use of the sunspot activity during the second half of the 20th century does not, on its own, necessarily indicate that solar activity was higher than previous periods, as the IAR asserts. Irradiance variations are driven not only by sunspots but also by facular and network magnetic features. Additionally, direct facular observations only extend back to 1892, and their use prior to the 1970s has been rather limited. Consequently, reconstructions of irradiance variations rely on empirical relationships between the sunspot number and facular and network irradiance uncertainty into estimates of the secular trend. However, uncertainties in these relationships, such as those in the observations (e.g., Chatzistergos et al. (2020)) and in revised irradiance reconstructions (e.g., Yeroshi, Chatzistergos et al. (2023)), can lead to a range of possible solutions with some of the reconstructions such as SATIRE-S, as mentioned above, even updated reconstructions with consistent isotope data (Wu et al. (2018)) suggest a small secular trend. Even when the correct is provided in the IAR report, standing in a potentially oversimplified or overstated implication that a large secular trend in solar irradiance, despite the balance of current evidence.

Quote given to Carbon Brief

Supporting evidence

- Long-term changes in solar activity and irradiance - ScienceDirect
- Reconstructions of the Sunspot Number: Status Report - Solar Physics
- Reviews of the SATIRE-S irradiance reconstruction model of TSI: Effects of instrumental and SATIRE-S model changes | Astronomy & Astrophysics (A&A)
- Understanding the secular variability of solar irradiance: the potential of CLA-S observations | Journal of Space Weather and Space Climate

8.3 Attribution of global warming

Page 33

MISLEADING

However, some scientists have concluded that it is not possible to be confident of a multi-decadal trend in TSI (Krivova, 2011).

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

This statement is misleading because it refers to section 5 of Krivova's (2011), which focuses on direct measurements of solar irradiance only since 1976, whereas the IAR report in the previous sentence was discussing the grand solar maximum and long-term trends beyond the period covered by direct TSI measurements. Furthermore, the uncertainty in reconstructing the long-term trend from direct measurements and modelled reconstructions, is widely acknowledged in the scientific literature and was further explored in a paper published by Krivova and co-authors. Recent evidence and advances consistently indicate a relatively weak secular trend in total solar irradiance (TSI) variations.

Quote given to Carbon Brief

Supporting evidence

- Long-term changes in solar activity and irradiance - ScienceDirect
- Changes in the Total Solar Irradiance and Climate Effects | Journal of Space Weather and Space Climate

8.3 Attribution of global warming

Page 33

MISLEADING

This uncertainty causes some reconstructions of TSI from 1750 to have low variability (implying a very low impact of solar variations on global mean surface temperature), whereas others with high TSI variability can explain more than 50% of the temperature variability since pre-industrial times (Scafetta (2013), Stefani (2021)).

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

Neither of the two papers cited for the claim that "datasets with high TSI variability can explain more than 50% of the temperature variability since pre-industrial times" (Stefani (2021) did not even use any TSI series; instead, he performed a linear regression between the temperature record and the sunspot number). Furthermore, the paper by Scafetta (Figure 1)B compares the Grand Sunspot Temperature (GST) record and the TSI model by Hoyt and Schatten that the ACRIM TSI composite was not considered. It observed since 1700, which suggests that the major observed climatic oscillations are well explained and that the solar contribution to the global warming since 1900 is small.

This raises two major concerns. First, the statement in the report is incorrect since neither Scafetta (2013) nor Stefani (2021) demonstrates that TSI can account for "*more than 50% of the temperature variability since pre-industrial times*". Second, the conclusion by Scafetta (2013) relies on the Hoyt and Schatten (1993) TSI model after demonstrating that it is in better agreement with the GST record. This model has been shown to be highly unrealistic and is now widely discredited (Chatzistergos 2020a). As detailed in Chatzistergos (2020a), the Hoyt and Schatten model suffers from multiple arbitrary adjustments and fabricated data for one index spanning over a century, and it is based on the false assumption that the Group Sunspot Number (GSN) is a linear proxy of the sunspot area (SSN) and that the latter is a linear proxy of TSI. When updated underlying indices and direct TSI composites (e.g., PMOD, Matthes et al. (2017), ACRIM) are considered, Krivova et al. (2007) have demonstrated that the exaggerated TSI variation amplitudes by roughly a factor of five (Chatzistergos 2020a). The discrepancy is even greater when considering the ACRIM TSI composite, which Hoyt and Schatten's model is incompatible with. This seriously undermines the reliability of conclusions drawn from the Hoyt and Schatten model for attributing recent climate change to solar variability.

Therefore, the two given references do not corroborate the claim of the IAR report. This statement likely originates from Scafetta et al. (2023), but the same criticisms previously discussed regarding Scafetta et al. (2023) apply here as well. Scafetta et al. (2023) relies on the Hoyt and Schatten TSI reconstruction, which is incompatible with the direct TSI measurements and is widely discredited. As noted in previous sections, its use in this context undermines the credibility of their results. While there is legitimate uncertainty about the exact magnitude of long-term solar irradiance trends, recent studies (Yeroshi et al. 2025) and (2025) exaggerated the solar impact by cherry-picking outdated and superseded data that disagree significantly from TSI variations.

Quote given to Carbon Brief

Supporting evidence

- A Reassessment of Radiosonde Total Solar Irradiance Trends Versus Their Drivers Over the Last Three Decades and a Half | Journal of Geophysical Research: Atmospheres
- The Planetary Scale of the Sun – Part – XXII – Geophysical Research Letters – Wiley Online Library

8.3 Attribution of global warming

Page 33

MISLEADING

This statement is misleading and overstating. First, the choice of TSI satellite record does not substantially influence the estimated solar contribution to climate change. Second, the statement is misleading because it implies that the Hoyt and Schatten TSI reconstruction, that was at least once used, is supported by other TSI reconstructions. Third, such reconstructions used by Connolly et al. (2021) to attribute most warming to the sun are outdated, superseded, and misleading. Finally, the statement that the uncertainty regarding the long-term trend in solar irradiance, which naturally affects the precise estimate of solar attribution and global warming, all scientific uncertainty in this contribution is relatively small. The sentence in the IAR report, however, dramatically underestimates the uncertainty in this contribution.

However, it is also possible that the authors of the DoE report intended to support that the choice of TSI satellite records influences the reconstruction outcomes and thus the estimated solar contribution to global warming. This interpretation would be misleading, as explained earlier, and they believe that:

1) While the choice of reference TSI record affects proxy model results depending on the regression approach, this effect is often small. For example, Chatzistergos et al. (2020) demonstrated that the choice of TSI reference has a relatively small impact on estimates to minimum trends in their regression model. Similarly, Figure 5 in Chatzistergos et al. (2021) shows only minor differences when using various TSI references in their century-long reconstructions.

2) This effect is typically negligible in physics-based semi-empirical models. Various studies have confirmed the minimal impact in SATIRE reconstructions when changing the reference TSI series (e.g., Chatzistergos et al. 2021). Notably, for the latest version of SATIRE-S, Chatzistergos et al. (2022) explicitly state that "*the trend in the updated SATIRE-S TSI composite is independent of measured TSI*".

3) As mentioned earlier, the IAR report cited Velasco Herrera et al. (2015) instead of irradiance reconstruction models, which may have caused the confusion. Velasco Herrera et al. (2015) presented a machine learning extrapolation of TSI that depends on the chosen TSI reference series. Given the chaotic nature of solar activity, such limited-data extrapolations can naturally exhaustion (see Petrovay (2010)). Moreover, Velasco Herrera et al. 2015 predictions for 2011–21 have already been contradicted by direct TSI measurements collected since.

Quote given to Carbon Brief

Supporting evidence

- Long-term changes in solar activity and irradiance - ScienceDirect

8.3 Attribution of global warming

Page 33

FALSE

Svalgaard et al. (2005) suggest their 'dike' or solar influence on climate might exist for the Grand Maximum.

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

The reference is incorrect, there is no Scafetta et al. (2005). There is Scafetta et al. (2006). Although Scafetta (2005) claimed that the Svalgaard et al. (2005) provides a dike of attributing most global warming to the sun mostly based on the use of the Hoyt and Schatten TSI reconstruction. The Svalgaard et al. (2005) and Scafetta (2006) have been roundly rejected as fundamentally flawed (e.g., Lockwood (2007), Ineson et al. (2011) and the one mentioned by Chatzistergos (2020a). Egorova et al. (2018) and the one recommended by CMIP6. Since the Hoyt and Schatten (1993) TSI series was not updated since its publication, the Svalgaard et al. (2005) is linking it to the ACRIM TSI composite. Chatzistergos (2020a) demonstrated that the high variability imposed by Hoyt and Schatten (1993) is inconsistent with direct TSI measurements and that a variability magnitude approximately five times smaller is more appropriate.

Additionally, Chatzistergos (2020a) showed that the Hoyt and Schatten (1993) model conflicts particularly with the ACRIM composite by exhibiting opposite trends during the so-called ACRIM gap, making Svalgaard et al. (2005) claim of a dike between TSI and ACRIM invalid. Thus, with Egorova et al. (2018) exceeding the Yeo et al. (2020) TSI composite, the Svalgaard et al. (2005) claim of a dike between TSI and ACRIM and the Hoyt and Schatten (1993) and Egorova et al. (2018) TSI models is invalid. Furthermore, Scafetta (2013) cites Svalgaard et al. (2005), Scafetta himself acknowledged that even when allowing for solar effects beyond irradiance, the Hoyt and Schatten model is inconsistent with the CMIP6 TSI forcing. Therefore, the claim that solar influence accounts for more than 50% of warming is unsupported.

Quote given to Carbon Brief

Supporting evidence

- A Reassessment of Radiosonde Total Solar Irradiance Trends Versus Their Drivers Over the Last Three Decades and a Half | Journal of Geophysical Research: Atmospheres

8.3 Attribution of global warming

Page 33

FALSE

There are numerous candidate processes, including solar ultraviolet changes, energetic particle precipitation, atmospheric electric field effect on cloud cover, cloud changes produced by solar modulation of galactic cosmic rays, large volcanic eruptions in the 20th Cent, and the strength of the solar wind. Each solar-induced effects are not included in current models, although indirect methods have been proposed (Lockwood and Fröhlich, 2008).

Dr Theodoros Christoudias, astrophysicist, Max Planck Institute for Solar System Research.

It is exaggerated and incorrect to claim that the impacts of such effects are significant. Many of these effects are indeed thought to be small in the IAR report, making their implications unclear (for example, "large volcanic changes in the magnetic field"). Furthermore, the statement that "*each solar-induced effects are not included in current models*" is misleading. As the sole highlights, as demonstrated by the CLOUD experiment at CERN (Pierce (2017)). While some effects on Earth's atmosphere from solar processes have been studied (e.g., Lockwood and Fröhlich (2008), Lockwood and Fröhlich (2020)), their effects are generally considered to be small.

Quote given to Carbon Brief

Supporting evidence

- Cosmic rays, aerosols, clouds, and climate: Recent findings from the CLOUD experiment - Pierce - 2017 - Reviews of Geophysics - Wiley Online Library
- Planetary Periodic Influences on the Sun's Variability - ADS
- Tentative electron precipitation into the atmosphere – ScienceDirect

8.3 Attribution of global warming

Page 33

MISLEADING

However, the recommended forcing dataset for the CMIP6 climate model simulations only includes the solar contribution variations since 1850 and does not include variations before 1850.

Prof Mike Lockwood, professor of space environment physics and president of the Royal Astronomical Society, University of Reading.

The statement is true, but very misleading as it is made to sound that larger TSI still pre-dated were ignored. There is only one long TSI with reconstruction and it is very much sub-rather – other reconstructions show much smaller drift. There is also evidence that some solar cycles that show that the TSI change reconstructions are correct.

Quote given to Carbon Brief

Supporting evidence

- Proxies, biases, cross-cross examinations to paint the irradiance and their contribution to total solar irradiance and solar radiative forcing of climate | Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences

8.3 Attribution of global warming

Page 33

MISLEADING

However, the impact of solar variations on the climate is uncertain and subject to substantial debate (Lockwood (2012), Lockwood et al. (2012), Lockwood et al. (2014), Lockwood et al. (2016), Lockwood et al. (2018)).

Prof Mike Lockwood, professor of space environment physics and president of the Royal Astronomical Society, University of Reading.

This is highly misleading as it conflates regional and global climate. The cited references were discussing regional impacts of solar variations where there is more uncertainty and some (limited) solar influence through decreased propagation of stratospheric responses, which are not relevant to the global mean surface temperature. Also, the solar influence is found to be reversing of the global stratosphere shows consistently, but the effect of solar variability on the climate is relatively small compared to effects of changing other climate influences.

Quote given to Carbon Brief

Supporting evidence

- Solar output and its influence on the changing thermal structure of the atmosphere | Weir

8.3 Attribution of global warming

Page 85

MISLEADING

However, the recommended forcing dataset for the CMIP6 climate model simulations used in AR6 for attribution studies averages two data sets with low solar variability (Matthes (2017).

Prof Katja Matthes, director, GEOMAR Helmholtz Centre for Ocean Research Kiel.

It is correct that the long-term trend in TSI is rather on the low variability side, but it is completely ignored that other more extreme trends (high variability datasets) are completely unrealistic. We only provide an extreme Maunder Minimum case as a sensitivity experiment.

Quote given to Carbon Brief

Supporting evidence

1. GMD - Solar forcing for CMIP6 (v3.2)

8.3 Attribution of global warming

Page 85

MISLEADING

While AR6 shows a substantially greater solar impact than does AR5, the overall impact of solar forcing on the climate was still assessed to be small compared to anthropogenic forcing. However, the impact ofsolar variations on the climate is uncertain and subject to substantial debate (Lockwood (2012); Connolly et al. (2021) – something that is not evident in the IPCC assessment reports. The variations of TSI over time remain a challenging problem. Since 1978, there have been directmeasurements of TSI from satellites. However, the data exhibits non-negligible inconsistencies, and interpreting any multi-decadal trends in TSI requires comparisons of observations from overlapping satellites. There are several rival composite TSI datasets that disagree as to whether TSI increased ordecreased during the period 1986-96 (the ACRIM gap; see Chapter 4). Further, the satellite record of TSI is used to calibrate proxy models that infer past solar variations from sunspots and cosmogenic isotope measurements (Velasco Herrera et al. (2015)).There is substantial evidence for high solar activity in the second half of the 20th century (starting in 1959) and extending into the 1990s, before a decline in the early 21st century; this period is often termedthe "Modern Maximum". (Chatzistergos et al. (2023); Solanki et al. (2004); Usoskin et al. (2007)). However, some scientists have concluded that it is not possible to be confident of any multi-decadal trend in TSI (Schmutz (2021)). This uncertainty causes some reconstructions of TSI from 1750 to have low variability (implying a very low impact of solar variations on global mean surface temperature) whereas datasets with high TSI variability can explain more than 70% of the temperature variability since pre-industrial times (Scafetta (2013); Stefani (2021)). The choice of TSI satellite record used in an analysis can therefore substantially influence how much climate change is attributed to human versus natural forcings.

Prof Katja Matthes, director, GEOMAR Helmholtz Centre for Ocean Research Kiel.

In Matthes et al. (2017), we conclude the following two points (just as an example of mis-citation):

1) A new and lower TSI value is recommended: the contemporary solar-cycle average is now 1,361.0 +/-0.5W/m^2 (Prša et al. (2016)).

2) Over the last three solar cycles in the satellite era, there is a slight negative TSI trend in the CMIP6 dataset. A recent reconstruction of the TSI, with a proper estimation of its uncertainties, suggests that this downward trend between the solar minima of 1986 and 2009 is not statistically significant (Dudok de Wit et al. (2017)). The TSI trend leads to an estimated radiative forcing on a global scale of -0.04 W/m^2, which is small in comparison with other forcings over this period.

Therefore the sentence in the DoE report stating that the "choice of TSI satellite record used in an analysis can therefore substantially influence how much climate change is attributed to human versus natural forcings" is completely wrong and ignores the thorough science. The radiative forcing to this natural forcing – even if we would have chosen a different TSI dataset – is small (-0.04 W/m^2).

The next completely wrong statement is that "such solar indirect effects are not included in climate models". In Matthes et al. (2017), we provide a comprehensive dataset including TSI and solar spectral irradiance data as well as solar particle forcing for the first time. We also provide an ozone dataset with the indirect effect included. Many of the CMIP6 models do include spectral solar irradiance as well as ozone changes, so the indirect effect and some even the particle effects. I am really shocked about these wrong statements. The IPCC is based on sound scientific evidence and this DoE report is kicking the evidence with feet. Misinterpreting and mis-citing or not at all citing important work (such as the important parts of Matthes et al. (2017) where we provide sound scientific evidence and discuss the details) – this is just horrible. By taking out only parts of the story and "simplifying it", it is becoming simply wrong. This is not sound science.

Quote given to Carbon Brief

Supporting evidence

1. GMD - Solar forcing for CMIP6 (v3.2)
2. NOMINAL VALUES FOR SELECTED SOLAR AND PLANETARY QUANTITIES: IAU 2015 RESOLUTION B3 - IOPscience
3. Methodology to create a new total solar irradiance record: Making a composite out of multiple data records - Dudok de Wit - 2017 - Geophysical Research Letters - Wiley Online Library

8.3 Attribution of global warming

Page 87

FALSE

The causes of the early 20th century warming are discussed by Hegerl et al. (2017 and 2019). The atmospheric CO_2 concentration increased from 298ppm in 1905 to 310ppm in 1941, implying that CO_2 had little impact. Volcanic activity during this period was very low and solar forcing is uncertain. Yet Hegerl et al. (2017) somehow inferred that 40-54% of this warming could be attributed to external forcing, with the rest associated with internal variability. Bronniman et al. (2024) focused on the causes of the cooling in the first decade of the 20th century in the southern hemisphere. They found that the cooling was related to a La-Niña-like pattern in the Pacific, a cold tropical and subtropical South Atlantic, a cold extratropical South Pacific, and cool southern midlatitude land areas. The Southern Annular Mode was positive, with a strengthened Amundsen–Bellingshausen seas low, although the spread of the data products is considerable. The warming in the 1930s and subsequent cooling during mid-century was particularly pronounced in the Arctic. Bokuchava and Semenov (2021) find that these variations were most likely caused by acombined effect of long-term natural climate variations in the North Atlantic and North Pacific with a contribution of the natural radiative forcing related to the reduced volcanic activity and variations of solar activity, as well as growing greenhouse gases concentrations. Tokinaga et al. (2017) showed that the combined effect of internally generated Pacific and Atlantic interdecadal variability intensified Arctic warming in the early 20th century. The synchronised Pacific-Atlantic warming drastically alters planetary-scale circulations over the Northern Hemisphere; these same circulation patterns have a global influence.

Dr Gabriele Hegerl, professor of climate system science, University of Edinburgh.

The authors discuss my papers on the early 20th century warming (Hegerl et al. (2017) and (2019)). For example, they argue that "the atmospheric CO_2 concentration increased from 298ppm in 1905 to 310ppm in 1941, implying that CO_2 had little impact". This is a faulty argument. First of, the analysis in the paper focuses on 1901-50, not the shorter period, and secondly, it cannot be argued that 12ppm has "little impact". It does have an impact, even though smaller than the total warming at the time (as shown in the paper). Indeed, it causes a noticeable warming of about 0.2C globally, based on our analysis (partly offset by aerosol cooling at the same time as shown in the paper). They also argue that volcanic activity was low – which is correct and exactly the reason for a small positive forcing as the climate moves towards a less volcanically forced background state (which we also found contributed roughly 0.1C). We really disagree with the characterisation of having "somehow" inferred that about half the warming was forced as this is shown and discussed in the paper as originating from a published Bayesian attribution analysis that accounts for model uncertainty - which is very different from an ad hoc "somehow" approach. The authors then discuss the important contribution unusual circulation made to this warming particularly in the Arctic, which we fully agree with. Regional short term variations are to a large extent driven by interesting variability, while a global mean trend over 50 years as in that case is driven to a substantial part by forcing. This is how climate works. Meanwhile, there are publications suggesting that some of the early 20th century cool start into the warming may be affected by data uncertainty – something an up to date DOE report maybe should have recognised and cited?

Quote given to Carbon Brief

Supporting evidence

1. The early 20th century warming: Anomalies, causes, and consequences - PubMed
2. Causes of climate change over the historical record - IOPscience
3. Early-twentieth-century cold bias in ocean surface temperature observations | Nature
4. Evaluating the Performance of Past Climate Model Projections - Hausfather - 2020 - Geophysical Research Letters - Wiley Online Library

8.3 Attribution of global warming

Page 88

MISLEADING

When the Great Pacific Climate Shift is accounted for in climate attribution analyses since 1950, 40% or more of the warming in the second half of the 20th century is attributed to natural internal variability (McLean et al. (2009); Tung and Zhou (2013); Chylek et al. (2016); and Scafetta (2021)).

Prof June-Yi Lee, professor of climate science, Pusan National University.

There are three problematic aspects in this contested text. First, the argument relies exclusively on Tung and Zhou (2013) and is not fully supported by the other three cited references. Specifically, McLean et al. (2009) and Scafetta (2021) provide no quantitative estimates of the contribution of internal variability to warming trend. Moreover, the estimate presented in Chylek et al. (2016) is substantially lower than that of Tung and Zhou (2013). According to Chylek et al. (2016), the contribution of the Atlantic Multidecadal Oscillation (AMO) to the 1970-2005 warming was between 0.13 and 0.20C (depending on the AMO index used), compared with a greenhouse gas contribution of 0.49-0.58C.

Second, while Figure 4 of Tung and Zhou (2013) suggests that internal variability may account for approximately 40% of the observed global warming over the past 50 years, it does not contribute significantly to the long-term warming trend over periods exceeding 75 years. This interpretation is consistent with the AR6 assessment (Section C.1, WGI SPM), which states: "Natural drivers and internal variability will modulate human-caused changes, especially at regional scales and in the near term, with little effect on centennial global warming. These modulations are important to consider in planning for the full range of possible changes."

Third, the statement does not acknowledge the ongoing debate regarding the role of external forcings in driving AMO variability. Recent studies have highlighted that what has traditionally been interpreted as an internally generated oscillation may, at least in part, reflect externally forced variability, thereby complicating to the estimation of the contribution of internal variability to global warming when relying solely on statistical analyses of observational data.

Quote given to Carbon Brief

Supporting evidence

1. Using data to attribute episodes of warming and cooling in instrumental records
2. The role of Atlantic Multi-decadal Oscillation in the global mean temperature variability | Climate Dynamics

8.3 Attribution of global warming

Page 89

MISLEADING

Section 8.3.3

Prof Andreas Schmittner and Prof Peter U. Clark, college of Earth, ocean and atmospheric science, Oregon State University.

In the second paragraph, the authors write: "An application to the well-known Vostok ice core data revealed an error in Al Gore's documentary An Inconvenient Truth. Gore showed the data and drew attention to the coherence of temperature and CO_2 changes over a 440,000 year span, which he asserted was due to CO_2 driving temperature changes. But temperature changes can also affect atmospheric CO_2 levels. Davidson et al. (2015) examined the series and found that temperature Granger causes CO_2 but not the reverse. In other words on the time scales represented in the Vostok data, the coherence in the series is primarily due to the influence of temperature on CO_2 levels, not the feedback of CO_2 levels on temperature." This section should clarify that the Vostok ice core data are oxygen isotopes, a proxy for regional temperature in Antarctica. They do not represent global temperature. Using a reconstruction of global temperature over the last deglaciation (last 20,000 years) Shakun et al. (2012) find that CO_2 leads temperature and thus likely causes much of the warming over that time period. This should be mentioned here.

This section is also contrary to the vast majority of palaeoclimate literature, which has established that atmospheric CO_2 has an important influence on global temperatures during past ice-age cycles. For example, radiative forcing of CO_2 and ice sheets similarly contribute to colder temperatures during the Last Glacial Maximum (Broccoli (2000); Broccoli & Manabe (1987); Hansen et al. (1984); Köhler et al. (2010); Schmittner et al. (2011)); and Weaver et al. (1998)) and simulations of glacial cycles with constant CO_2 underestimate ice volume changes (Abe-Ouchi et al. (2013) and Ganopolski & Calov (2011)). This should be mentioned here, since otherwise it will leave the reader with the false impression that CO_2 variations are unimportant for ice age climate changes.

In the first paragraph, there are several non-peer reviewed citations – e.g. Razzak (2021) and Dagsvik and Moen (2023). The Razzak (2021) paper is problematic because it removes the trends in both CO_2 and temperature data and then investigates the residuals, finding no correlations using only statistical methods. However, physical understanding of Earth's radiation budget clearly suggests that the trend in CO_2 causes the observed temperature trend (Pierrehumbert (2011)). Thus, removing the trends in both time series is not appropriate for attribution studies. Razzak (2021) also ignores other external forcings such as aerosols. I suggest to remove Razzak (2021) and other non-peer reviewed citations. Studies that include other forcings find convincing effects of CO_2 on global warming over the last 100 years (e.g. Lean and Rind (2009)).

Quote given to Carbon Brief

Supporting evidence

1. Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume | Nature
2. Tropical Cooling at the Last Glacial Maximum: An Atmosphere–Mixed Layer Ocean Model Simulation in: Journal of Climate Volume 13 Issue 5 (2000)
3. The influence of continental ice, atmospheric CO2, and land albedo on the climate of the last glacial maximum | Climate Dynamics
4. CP - The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles
5. Climate Sensitivity: Analysis of Feedback Mechanisms - Hansen - 1984 - Geophysical Monograph Series - Wiley Online Library
6. What caused Earth's temperature variations during the last 800,000 years? Data-based evidence on radiative forcing and constraints on climate sensitivity - ScienceDirect
7. How will Earth's surface temperature change in future decades? - Lean - 2009 - Geophysical Research Letters - Wiley Online Library
8. Infrared radiation and planetary temperature | Physics Today | AIP Publishing
9. Climate Sensitivity Estimated from Temperature Reconstructions of the Last Glacial Maximum | Science
10. Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation | Nature
11. Simulated influence of carbon dioxide, orbital forcing and ice sheets on the climate of the Last Glacial Maximum | Nature

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

Arguably the most striking change in the Earth's climate system during the 21st century is a significant reduction in planetary albedo since 2015, which has coincided with at least two years of record global warmth. Figure 8.2 shows the planetary albedo variations since 2000, when there are good satellite observations. The 0.5% reduction in planetary albedo since 2015 corresponds to an increase of 1.7 W/m^2 in absorbed solar radiation averaged over the planet (Hansen and Karecha (2025)). For comparison, Forster et al. (2024) estimate the current forcing from the increase in atmospheric CO_2 compared to pre-industrial times to be 2.33 W/m^2.

Prof Piers Forster, professor of climate physics, University of Leeds.

[The citation] is correct, but it presents a somewhat incomplete comparison with the absorbed solar radiation. One's a very long-term forcing, the other is a mix of forcing and response over a shorter period, so the two numbers are not very comparable. All Earth's energy budget terms need to be considered and understood together. And when all components are analysed together, the cause of the Earth's energy balance can be traced to GHG warming and the Earth's system response with a good degree of certainty, as explained by the Hodnebrog et al. (2024) paper already referenced. This text as written is ok, but for me it plays up the uncertainties somewhat.

Quote given to Carbon Brief

Supporting evidence

1. ESSD - Indicators of Global Climate Change 2024: annual update of key indicators of the state of the climate system and human influence
2. Recent reductions in aerosol emissions have increased Earth's energy imbalance | Communications Earth & Environment

8.4 Declining planetary albedo and recent record warmth

Page 90

FALSE

Arctic sea ice extent has declined by about 5% since 1980

Anonymous.

September sea ice (1979-88 average compared to 2015-24 average) is 34% lower today according to OSISAF [Ocean and Sea Ice Satellite Application Facility] observations and 35% lower according to NSIDC [National Snow and Ice Data Center]. For the annual mean sea ice extent, the percentage reduction is 14% according to NSIDC and OSISAF. The number is higher in summer because that is when there is the most sea ice loss and the seasonal minimum, so the least amount to start with. In addition, the evidence given to support this statement links to a figure showing sea ice decline in Antarctica in July.

Quote given to Carbon Brief

Supporting evidence

1. Sea-ice index | OSI SAF
2. Data Tools | National Snow and Ice Data Center

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

...although following 2007 there has been a pause in the Arctic sea ice decline (England et al. (2025)).

Anonymous.

What they say is true and is correctly cited, but lacking context. In the latest version published recently [the DoE report cites a preprint version], the authors note that they "find that the existence of this slowdown also predisposes the sea ice cover for a more rapid decline in the near future". They add that they "would like to underscore that pause or slowdown are used interchangeably to refer to an extended period with little or no decline in sea ice cover, due to the observed realisation of multi-decadal climate variability on top of the response to anthropogenic forcing, temporarily interrupting the ongoing long-term reduction in Arctic sea ice. This does not imply a cessation of human-induced climate change and, instead, it is likely that sea ice would have increased over this period without human influence."

Quote given to Carbon Brief

Supporting evidence

1. Minimal Arctic Sea Ice Loss in the Last 20 Years, Consistent With Internal Climate Variability - England - 2025 - Geophysical Research Letters - Wiley Online Library

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

Regarding Antarctic sea ice, the IPCC AR6 concludes that "There has been no significant trend in Antarctic sea ice area from 1979 to 2020 due to regionally opposing trends and large internal variability." (Summary for Policymakers, A.1.5)

Anonymous.

While this is true, it is misleading because they neglect to mention that Antarctic sea ice cover has had a dramatic decline since 2015 and the last few years have seen record lows. Studies have found that these record lows were unlikely to have happened before over the last century. Missing out the last few years of Antarctic sea ice trajectory is a huge omission.

Quote given to Carbon Brief

Supporting evidence

1. A regime shift in seasonal total Antarctic sea ice extent in the twentieth century | Nature Climate Change
2. A twenty-first century structural change in Antarctica's sea ice system | Communications Earth & Environment

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

Section 8.4

Dr Helge F. Goessling, climate physicist and group leader, Alfred Wegener Institute.

The authors manage to avoid statements that are individually clearly wrong. It is even somewhat difficult to nail down individual sentences that would be clearly misleading taken alone, because that depends on the context they are placed in. However, considering the section as a whole, I think it's pretty clear that they are over-emphasising uncertainties and the possible relative role of natural variability while down-playing the two other main possible mechanisms behind the albedo decline of the last two and a half decades, namely a possibly emerging positive low-cloud feedback and (largely indirect) aerosol effects.

Quote given to Carbon Brief

Supporting evidence

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

Section 8.4

Prof Richard Allan, professor of climate science, University of Reading.

I did not spot major errors in the section on planetary albedo, although I consider that there is an emphasis on abruptness of temperature and albedo changes, which artificially emphasise the role of natural variability. which could be misleading.

Quote given to Carbon Brief

Supporting evidence

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

A sharp recent increase in global average temperatures has raised the question of short-term drivers of climate. One such candidate is the fraction of absorbed solar radiation which has also increased abruptly in recent years. The question is whether the change is an internal feedback to warming caused by greenhouse gases, or whether something else increased the fraction of absorbed radiation which then caused the recent warming.

Prof Richard Allan, professor of climate science, University of Reading.

A "sharp recent increase in global temperature" and "the fraction of absorbed solar radiation which has also increased abruptly" are potentially misleading since while both global surface temperature and absorbed solar radiation have varied with ENSO, they have also increased steadily over time (Loeb et al. (2024) and Forster et al. (2024)). This wording could artificially emphasise the role of natural variability.

Quote given to Carbon Brief

Supporting evidence

1. Observational Assessment of Changes in Earth's Energy Imbalance Since 2000 | Surveys in Geophysics

8.4 Declining planetary albedo and recent record warmth

Page 90

MISLEADING

Arguably the most striking change in the Earth's climate system during the 21st century is a significant reduction in planetary albedo since 2015, which has coincided with at least two years of record global warmth.

Prof Richard Allan, professor of climate science, University of Reading.

Planetary albedo has decreased before 2015 as well as after, though the decreases are larger around 2012-14 (increases in absorbed sunlight in Loeb et al. (2024), Fig. 5a). Decreases in reflected sunlight up to 2016 are captured by some climate models applying observed SST (Loeb et al. (2020)), suggesting that they are a response to global warming and its spatial pattern (Andrews et al. (2022)), though it is not clear how much of the pattern of global warming is explained by radiative forcing and how much internal variability.

Quote given to Carbon Brief

Supporting evidence

1. New Generation of Climate Models Track Recent Unprecedented Changes in Earth's Radiation Budget Observed by CERES - Loeb - 2020 - Geophysical Research Letters - Wiley Online Library

8.4 Declining planetary albedo and recent record warmth

Page 91

MISLEADING

The issue then becomes the cause of the change in cloud cover. Two explanations have been posited for the declining cloud cover over the past decade: • Natural climate variability • Changes in low cloud cover associated with warming sea surface temperatures, implying an emerging positive feedback to climate change (Hansen and Karecha (2025))

Dr Helge F. Goessling, climate physicist and group leader, Alfred Wegener Institute.

Maybe even more importantly, they omit the third major possible mechanism behind the cloud-cover decline altogether, namely indirect aerosol effects. They do mention aerosols and recent aerosol emission reductions, but only prior to the part where they state that the main reason for the albedo-decline seems to be related to clouds (which, by itself, is true). They write: "The issue then becomes the cause of the change in cloud cover. Two explanations have been posited for the declining cloud cover over the past decade." However, the indirect aerosol effect, although uncertain, is likely even stronger than the direct aerosol effect, and it is an important candidate for explaining a part of the recent cloud-cover decline. By omitting this completely, they further emphasise the relative importance of natural variability to a degree that can not be considered appropriate.

Quote given to Carbon Brief

Supporting evidence

8.4 Declining planetary albedo and recent record warmth

Page 91

MISLEADING

Surface albedo changes have thus contributed only weakly to the recent planetary albedo decline, particularly when averaged annually and globally.

Prof Richard Allan, professor of climate science, University of Reading.

Surface albedo has also contributed significantly to the decline in global albedo according to Loeb et al. (2024), which accounts for cloud masking: "...part of the positive –SW trend is impacted by decreases in surface albedo from declining sea-ice coverage during the CERES period". However, this applies to the period since 2000, rather than during 2015, which line 10 [of the DoE report] is referring to, while the decrease in planetary albedo has been observed over the full CERES period rather than just in 2015 (Loeb et al. (2024)).

Quote given to Carbon Brief

Supporting evidence

1. Observational Assessment of Changes in Earth's Energy Imbalance Since 2000 | Surveys in Geophysics

8.4 Declining planetary albedo and recent record warmth

Page 91

MISLEADING

Loeb et al. found that decreases in low- and mid-level clouds since 2015 are the primary reason for decreasing planetary albedo in the northern hemisphere, whereas in the southern hemisphere the decrease in planetary albedo is primarily due to decreases in mid-level clouds across all latitude zones.

Prof Richard Allan, professor of climate science, University of Reading.

The stated "decreases in low- and mid-level clouds" is misleading since Loeb et al. (2024) find decreases in cloud cover and also reflectance have contributed to the increase in absorbed sunlight along with surface albedo decreases (that can are related to ice melt elsewhere in the article): "[W]e find that decreases in low and middle cloud fraction and reflection and reduced reflection from cloud-free areas in mid-high latitudes are the primary reasons for increasing ASR [absorbed solar radiation] trends in the [northern hemisphere]...In the [southern hemisphere] the increase in ASR is primarily from decreases in middle cloud reflection and a weaker reduction in low-cloud reflection." The increase in reflection counters arguments presented that the decreases in aerosol are not influencing the decreases in albedo as their effect on making clouds brighter diminishes (e.g. Hodnebrog et al. (2024)), while the additional steady rise in ASR during the CERES period is also consistent with cloud and ice-albedo feedbacks to warming (Forster et al. (2021); Tselioudis et al. (2025); and Norris et al. (2016)) as well as declining global aerosol emissions (Quaas et al. (2022)).

Quote given to Carbon Brief

Supporting evidence

1. Observational Assessment of Changes in Earth's Energy Imbalance Since 2000 | Surveys in Geophysics

8.4 Declining planetary albedo and recent record warmth

Page 91

MISLEADING

The issue then becomes the cause of the change in cloud cover. Two explanations have been posited for the declining cloud cover over the past decade: • Natural climate variability • Changes in low cloud cover associated with warming sea surface temperatures, implying an emerging positive feedback to climate change (Hansen and Karecha (2025))

Prof Richard Allan, professor of climate science, University of Reading.

Two explanations for decreased global albedo are presented, but the decline in ocean aerosols is ignored and this has been identified as an important driver of decreased global albedo (e.g. Hodnebroeg et al. (2024)). Combined with overemphasis of the abruptness of changes, this is potentially misleading since it overemphasises the role of natural variability.

Quote given to Carbon Brief

Supporting evidence

1. Recent reductions in aerosol emissions have increased Earth's energy imbalance | Communications Earth & Environment

8.4 Declining planetary albedo and recent record warmth

Page 92

MISLEADING

It is not easy to justify a new positive low cloud feedback that began emerging in 2015 since there is no obvious feedback trigger starting at that time.

Dr Helge F. Goessling, climate physicist and group leader, Alfred Wegener Institute.

I think this is an example of a "straw man argument", because it's clear that there would not be a trigger starting exactly at some point. Rather, a forced signal (like a positive cloud feedback) would be superimposed by natural variations, and at some point, depending on signal-to-noise ratio (and possible other forcings) and how the associated forcing grows, the forced signal can emerge from the noise. An associated timeseries may just look like something kicked in at a specific time.

Quote given to Carbon Brief

Supporting evidence

8.4 Declining planetary albedo and recent record warmth

Page 92

MISLEADING

However, there are numerous natural climate signals during this period that are associated with atmospheric circulation changes that can influence the distribution of clouds: • The 2014-16 was one of the strongest El Niño events on record. • A cold anomaly beginning in 2015 in the subpolar gyre of the North Atlantic reflects a shift in the ocean circulation pattern associated with decadal variability in the Atlantic (Frajka-Williams et al. (2017); Arthur et al. (2021)). • The Pacific Decadal Oscillation positive index peaked in 2016, then declined and has been in negative territory since late 2019. • Eruption of the submarine Hunga-Tonga volcano in 2022.

Dr Helge F. Goessling, climate physicist and group leader, Alfred Wegener Institute.

When it comes to natural variability, they are listing quite comprehensively which aspects of natural variations could – in principle – have had an impact on planetary albedo, in part also overemphasising some aspects that rather clearly can not have had a major impact on the longer-term decline, such as ENSO (as we also investigated in our paper). Also their interpretation of results for the Hunga Tonga eruption and its possible contribution to the recent warming surge seems not to be a balanced representation of the recent literature about it, which does not suggest a major contribution. Also, they provide their list of natural variability aspects right after the down-playing statement about the possible low-cloud feedback mentioned above, saying: "However, there are numerous natural climate signals during this period that are associated with atmospheric circulation changes that can influence the distribution of clouds." This shows quite clearly that they want to emphasise the natural variability over the possible low-cloud feedback.

Quote given to Carbon Brief

Supporting evidence

8.4 Declining planetary albedo and recent record warmth

Page 92

MISLEADING

It is not easy to justify a new positive low cloud feedback that began emerging in 2015 since there is no obvious feedback trigger starting at that time.

Prof Richard Allan, professor of climate science, University of Reading.

A change in the SST pattern has been linked to a decline in low-altitude cloud over some stratocumulus regions (Andrews et al. (2022) and Loeb et al. (2020)) and it is misleading to paint these changes as requiring new feedbacks.

Quote given to Carbon Brief

Supporting evidence

1. New Generation of Climate Models Track Recent Unprecedented Changes in Earth's Radiation Budget Observed by CERES - Loeb - 2020 - Geophysical Research Letters - Wiley Online Library

8.4 Declining planetary albedo and recent record warmth

Page 92

MISLEADING

A change of 1-2% in global cloud cover has a greater radiative impact on the climate than the direct radiative effect of doubling CO2.

Prof Richard Allan, professor of climate science, University of Reading.

Short-term interannual fluctuations are unfairly compared with long-term radiative forcing. Also, a decrease in global albedo is an expected consequence of global warming as feedbacks cause ice to melt and some clouds to diminish (e.g. Forster et al. (2021)). This signal in shortwave radiation is seen to a greater extent than in the longwave since decreases in outgoing longwave due to rising greenhouse gases being offset by the increases in outgoing longwave from the resulting higher temperatures that are influenced by the greenhouse gases, but also declining aerosol and feedbacks that amplify the warming through ice melt and cloud responses to warming (e.g. Raghuraman et al. (2024)).

Quote given to Carbon Brief

Supporting evidence

1. Chapter 7: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity | Climate Change 2021: The Physical Science Basis

8.6 Extreme event attribution (EEA)

Page 96

FALSE

The existence of an outlier at the end of a data series poses the problem that estimates of the event probabilities will be biased whether the outlier is included or excluded (Barlow et al. (2020)). Methods to eliminate the bias have not yet been established.

Dr Ophélia Miralles, research scientist, ETH Zurich and Prof Anthony C. Davison, professor of mathematics, École Polytechnique Fédérale de Lausanne.

The text suggests that the presence of the trigger event at the end can't be dealt with, whereas the argument in Barlow and in our paper is that it can be dealt with (if the selection event is known) with appropriate statistical methods specifically detailed in both papers.

Quote given to Carbon Brief

Supporting evidence

1. Timing and spatial selection bias in rapid extreme event attribution - ScienceDirect
2. Inference for Extreme Values Under Threshold-Based Stopping Rules | Journal of the Royal Statistical Society Series C: Applied Statistics | Oxford Academic

8.6 Extreme event attribution (EEA)

Page 96

MISLEADING

Methods to eliminate the bias have not yet been established, leading some experts (e.g. Miralles and Davison (2023)) to argue that in settings in which a data series contains a single extreme event at the end, estimation of a return period for the extreme event will be so biased and uncertain that it should be avoided altogether.

Dr Ophélia Miralles, research scientist, ETH Zurich and Prof Anthony C. Davison, professor of mathematics, École Polytechnique Fédérale de Lausanne.

The cited paper does argue that the estimated return period is highly variable and so one should avoid using it, but also states that if it has to be used, one should clearly state its uncertainty. It does not imply that fast weather attribution should not be performed at all.

Quote given to Carbon Brief

Supporting evidence

1. Timing and spatial selection bias in rapid extreme event attribution - ScienceDirect

8.6 Extreme event attribution (EEA)

Page 97

MISLEADING

Zeder et al. (2023) also concluded that the methods employed by Philip et al. (2022, the WWA analysis) tend to overstate the rarity of extreme heat waves, leading to a biased perception of the effect of climate change on the heatwave event: "The tendency to overestimate the return period of observed extreme heatwave events may fuel the impression that seemingly impossible heatwave extremes are currently clustering at an unprecedented rate."

Prof Erich Fischer, lecturer at the department of environmental systems science, ETH Zurich.

The quote taken from the paper Zeder et al. (2023) led by my (former) PhD student, Joel Zeder, is in itself correct, yet taken completely out of context. Our paper shows indeed that when using relatively short observational timeseries, return level estimates may be systematically underestimated and return periods may be overestimated, which affects the calculation of the risk ratio in attribution studies. We further concluded that the risk of extreme heatwaves could be underestimated in both past and present-day climate and recommend following an approach like Miralles and Davison (2023). This paper applied the recommended approach specifically to the 2021 Pacific north-west heatwave. However, both Miralles and Davison (2023) and Zeder et al. (2023) show that while methodological choices affect the exact estimate of the risk ratio, all estimates show a strong role of long-term warming increasing the probability of the event.

Quote given to Carbon Brief

Supporting evidence

1. The Effect of a Short Observational Record on the Statistics of Temperature Extremes - Zeder - 2023 - Geophysical Research Letters - Wiley Online Library
2. Timing and spatial selection bias in rapid extreme event attribution - ScienceDirect

Part III: Impacts on ecosystems and society

9.1 Econometric analyses

Page 104

FALSE

Bareille and Chakir (2023) assembled a large data base on farm sale prices in France for properties that sold twice between 1996 and 2019. They could replicate pessimistic results showing negative effects of warming on agricultural land values using conventional econometric modelling.

Dr Francois Bareille, researcher in economics, INRAE & University Paris-Saclay.

Bareille and Chakir (2023) do reproduce the standard results obtained from "conventional econometric modelling" like the Ricardian approach proposed by Mendelsohn et al. (1994). However, these standard results in no way demonstrate "negative effects of warming on agricultural land values". Rather, as illustrated in Figure 5 of their study, Bareille and Chakir (2023) closely mimic the outcomes of the Ricardian approach in similar contexts, which points to slightly positive impacts of climate change on farmland values. Such findings are consistent with the broader literature when applied to contexts with moderate climate conditions. Nevertheless, as has been extensively discussed, these results are likely subject to bias arising from omitted variables.

Quote given to Carbon Brief

Supporting evidence

1. The impact of climate change on agriculture: A repeat-Ricardian analysis - ScienceDirect
2. The empirical analysis of climate change impacts and adaptation in agriculture - ScienceDirect
3. The Use of Cross-Sectional Analysis to Measure Climate Impacts on Agriculture: Theory and Evidence | Review of Environmental Economics and Policy: Vol 11, No 2

9.1 Econometric analyses

Page 104

FALSE

But by taking advantage of the repeat sales data, which provides information on site-specific changes in land prices, they found the results reversed and implied that climate change will be very beneficial for French agriculture.

Dr Francois Bareille, researcher in economics, INRAE & University Paris-Saclay.

Bareille and Chakir (2023) do not identify any negative impacts of climate change on land values; across all methodologies they employ, the evidence consistently points toward positive effects of recent climate trends.

Quote given to Carbon Brief

Supporting evidence

1. The impact of climate change on agriculture: A repeat-Ricardian analysis - ScienceDirect

9.1 Econometric analyses

Page 104

MISLEADING

But by taking advantage of the repeat sales data, which provides information on site-specific changes in land prices, they found the results reversed and implied that climate change will be very beneficial for French agriculture.

Dr Francois Bareille, researcher in economics, INRAE & University Paris-Saclay.

These results are derived from recent climate trends over the period 1990-2020 in the context of French agriculture. While such moderate warming may generate short-term benefits for agriculture, more substantial – and as yet unobserved – warming is likely to result in losses. It is well established that higher temperatures induced by climate change can produce non-linear effects, with moderate warming potentially yielding gains, but more pronounced warming leading to detrimental impacts.

Quote given to Carbon Brief

Supporting evidence

1. Global non-linear effect of temperature on economic production | Nature

9.1 Econometric analyses

Page 104

MISLEADING

But by taking advantage of the repeat sales data, which provides information on site-specific changes in land prices, they found the results reversed and implied that climate change will be very beneficial for French agriculture.

Dr Francois Bareille, researcher in economics, INRAE & University Paris-Saclay.

While such a result may arise in contexts like France (where northern regions remain, for the time being, too cold to cultivate high-value crops), these findings are highly context-specific and cannot be generalised to other settings. This is particularly true for a large and heterogeneous country like the US, where both climatic and agricultural conditions differ substantially from those in France. Bareille and Chakir (2023) provide indirect evidence that the positive impacts of recent climate change on French agriculture identified with their approach are likely driven by vineyard expansion, a phenomenon highly specific to France and not transferable to the US context.

Quote given to Carbon Brief

Supporting evidence

1. The Use of Cross-Sectional Analysis to Measure Climate Impacts on Agriculture: Theory and Evidence | Review of Environmental Economics and Policy: Vol 11, No 2

9.1 Econometric analyses

Page 104

MISLEADING

The authors concluded that, taking adaptation into account, a warming climate would yield positive benefits for French agriculture that were between two and 20 times larger than had previously been estimated. On average, with full adaptation, they concluded that climate change under the medium RCP4.5 scenario could double the value of French farmland by 2100

Dr Frances Bentley, researcher in taxonomy, HMAE & University Parry Sarkaj

These results are obtained using a novel method, the so-called repeat-fluctuation approach. In contrast to other studies cited in the report – such as Mookherjee et al. (2015), Stockdale and Constantinidou (2018), Lefebvre-Dubois (2019) and Burke and Emerick (2016) – this approach has not been replicated in other contexts. Its findings should therefore be treated as provisional. For a detailed discussion of the implications and extensions of Stockdale and Constantinidou (2018), see Lefebvre-Dubois (2019) and Burke and Emerick (2016) respectively. Further empirical research on impacts of climate change on crop productivity in the US, but also worldwide.

Return to table of contents

Supporting evidence

- Impacts of climate change on global agriculture accounting for adaptation - Report

Dr Frances Bentley, researcher in taxonomy, HMAE & University Parry Sarkaj

All of these studies implicitly account for the effects of CO2 fertilisation on agriculture, insofar as local CO2 concentrations are correlated with changes in temperature and precipitation. Moreover, Hallgrave et al. (2015) show that the positive effects of CO2 fertilisation only partially offset the negative impacts of anthropogenic climate change on crop yields arising from changes in temperature and precipitation.

Return to table of contents

Supporting evidence

- Impacts of climate change on global agriculture accounting for adaptation - Report

9.2 Field and laboratory studies of CO2 enrichment

Page 127

Field and laboratory studies of CO2 enrichment have been conducted through the use of [redacted] to expose crops to elevated CO2 concentrations. Typically, these studies find that [redacted] conditions are recorded. Ahmadi et al. (2020) summarise results from about 250 [redacted] studies. They find an average CO2 fertilisation effect of about [redacted] per cent, with a typical [redacted] per cent. Effects are found to be smaller in [redacted] studies that more closely resemble the natural climate system. Ideally, increasing the usable sample size by 40%. The gap yield projections incorporating the results of [redacted] studies are presented in the table shown in Figure 9.2, whereas the partial data set implied warming would decrease yields. Only then, the complete data set would be estimated to increase global yields, since the CO2 warming system keeps.

Dr Delphine Devoy, lead author, IPCC WG1 WG2.

Regression results and statistical climate impacts on crop yield do not capture the impact of non-linear processes, such as effects of extreme heat stress occurring at crop flowering. Schlenker and Roberts (2009) find that yields for maize, soybeans, and cotton fall by 50% at 32°C, 34°C, and 36°C, respectively. Temperatures above these thresholds are common in the US, but also worldwide. Ignoring these effects will lead to an overestimate of the impact of climate change on crop yields, as temperatures increase.

Return to table of contents

Supporting evidence

- Chamber, De Bondt, Hemmerly, Climate Change 2022: Impacts, Adaptation and Vulnerability

9.3 Cop modelling meta-analyses

Page 128

Dr Delphine Devoy, lead author, IPCC WG1 WG2.

Regression results and statistical climate impacts on crop yield do not capture the impact of non-linear processes, such as effects of extreme heat stress occurring at crop flowering. Schlenker and Roberts (2009) find that yields for maize, soybeans, and cotton fall by 50% at 32°C, 34°C, and 36°C, respectively. These temperatures are common in the US, but also worldwide. Ignoring these effects will lead to an overestimate of the impact of climate change on crop yields, as temperatures increase.

Return to table of contents

Supporting evidence

- Chamber, De Bondt, Hemmerly, Climate Change 2022: Impacts, Adaptation and Vulnerability

9.4 CO2 fertilisation and nutrient loss

Page 127

Evidence has shown that CO2-induced biomass gains are sometimes accompanied by reductions in the concentrations of protein and other key nutrients such as iron and zinc [the ref. 2015].

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

All plants absorb carbon dioxide from the atmosphere, using photosynthesis pathways that differ in the number of carbon atoms in the molecules that are formed when carbon is first taken in by the plant. Crops that use the Calvin cycle, which includes important crops such as wheat, rice, barley, soy, and potatoes, are called C3 plants. Field experiments with wheat and rice have shown that when grown at the elevated carbon dioxide concentrations that are projected to occur later this century, the protein concentration of these C3 crops declines by about 10%. Protein concentration in wheat grain is reduced by 7.6% (with a 95% confidence interval of 6.4% to 8.8%) and in rice grain by 14.1% (12.4% to 15.8%). This observation is of particular concern because about 30% of calories consumed worldwide, with wide regional variability (Avila-Cabadilla et al. 2009). Some of the lost protein can be replaced in the diet, but the photosynthetic pathway (C4 or not) are expected to experience declines in nutrient concentrations. Free-air carbon dioxide enrichment (FACE) experiments with wheat, rice, and potatoes have found that when grown at 550 parts per million, the micronutrient density are consistently seen under elevated CO2 by 5% (3.2% to 7.1%), 5.1% (3.8% to 6.4%), 6.3% (4.9% to 7.7%), and 8.1% (6.5% to 9.7%), respectively. Similar results have been found for barley (4.8%), soybeans (10.8%), and rice (8.1%), which are widely consumed in the US.

The fact CMS report also ignores the fact a yield loss at 550 ppm CO2 is more than offset by elevated temperature can increase nutrient loss as well as, and possibly even more than, yield loss (Zhao et al. 2020; Fatima et al. 2021; 2022; Gao et al. 2011; Mueller et al. 2012; Myers et al. 2014; Smith et al. 2000; Tubiello et al. 2007; Zhu et al. 2018). Yet, it still has diluted nutrients. Ingestion of foods in food recalls in Illinois, shortened life span, increased risk of heart disease, and [redacted] (Ebi et al. 2021).

9.4 CO2 fertilisation and nutrient loss

Page 127

Some experiments have shown that the rising temperatures expected to accompany higher CO2 levels will offset this loss (Myers et al. 2014) although the effect is not fully understood, as the evidence that nutrient dilution observed to date is entirely attributable to higher CO2 (Pan 2020).

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

Some experiments have shown that rising temperatures expected to accompany higher CO2 levels can offset nutritional losses (Myers et al. 2019), and while the Dak DMS project acknowledges that evidence for this compensation is not fully understood, it looks to the studies that found partial or no compensation in nutritional density when temperatures increased. Experimental evidence for the Dak DMS project is from Loladze (2014), Wang et al. (2019), Wei et al. (2012), and Zhu et al. (1997). On the flip side of crop nutrition, the experiments that found no compensation are from Loladze (2014) and Myers et al. (2014). Wang et al. (2020) found that while warmer temperatures moderately mitigated the loss of protein and micronutrients in rice grain, the net effect was still a decrease in micronutrient density and protein concentration in rice grain.

The second half of the statement above is "... as is the evidence that nutrient dilution observed to date is entirely attributable to higher CO2" (shown as incorrect by Loladze 2014; Myers 2015; Pan 2020) presents a figure (Figure 9.4) showing that warmer breeding lines of spring wheat have six percent protein, but all that 20 percent of the wheat breeding lines in present protein are in the 6% to 8% range. Thus, Dak's DMS project's statement that crop change nutritional density has "ultimately conclusive" [sic] is directly disproved by presents concentration separate from breeding factors.

Return to table of contents

Supporting evidence

- Increased temperature may safeguard the nutritional quality of crops under future elevated CO2 concentrations - Keller, 2017 - The Plant Journal - Wiley Online Library
- Effects of elevated CO2 and temperature on seed composition and phenology in plants from an alpine-subalpine natural grassland ecosystem - Gao - 2013 - Physiologia Plantarum - Wiley Online Library
- Changes in grain protein and amino acid concentrations of wheat and rice under short-term increased CO2 and temperature in isotope-labelled 15N experiments - Zhao - 2020 - Journal of Agronomy and Crop Science - Wiley Online Library
- Effects of elevated carbon dioxide and ozone on grain yield and quality of a high-yielding hybrid rice cultivar grown in China - Zhu - 2018 - Journal of the Science of Food and Agriculture - Wiley Online Library
- Effects of elevated [CO2] on grain nutrient concentrations in Fe-deficient rice depend on Fe supply - Myers - 2014 - New Phytologist - Wiley Online Library
- Responses of wheat and rice to post-anthesh warming in elevated carbon dioxide and canopy warming - Fatima - 2021
- Post-anthesis Heat Exacerbated Grain Nutrient Concentration Under Elevated CO2 Environment in Rice - Fatima - 2022 - Journal of the Science of Food and Agriculture - Wiley Online Library
- Rising Carbon Dioxide and Global Nutrition: Evidence and Action Needed

9.4 CO2 fertilisation and nutrient loss

Page 127

Evidence has shown that under rising CO2 levels, there are several adaptive responses that can occur in plants, such as a decline in breeding or leaf nitrogen content in a range of C3 and C4 species [the ref. 2015] and has proven to be a net offsetted agronomic strategy [the ref. 2015].

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

Biofortification via selective breeding has been used successfully to raise the zinc, iron and vitamin A content of crops, per Saltzman et al. (2017). However, this approach is not a substitute for the use of fertilisers, as the yield gains from biofortification are not as numerous nutrient thickness be elevated CO2. Biofortification is not a substitute for the use of increased levels in crops associated with elevated CO2 and CO2-induced warming.

Return to table of contents

Supporting evidence

- Availability, production, and consumption of crops biofortified by plant breeding: current evidence and perspectives - Saltzman - 2017 - Annals of the New York Academy of Sciences - Wiley Online Library
- Nutritional quality of crops in a future CO2 world: an agenda for research to mitigate food insecurity - Lobell - 2022 - Annual Review of Environment and Resources
- Effects of elevated CO2 on the nutritional quality of pollen and associated physiological impacts in plants - Singh - 2018 - Plant Physiology and Biochemistry
- Responses of wheat and rice to post-anthesis heat stress under different rates of leaf canopy nitrogen and atmospheric CO2 concentration - Fatima - 2020 - Agronomy - MDPI
- Post-anthesis Heat Exacerbated Grain Nutrient Concentration Under Elevated CO2 Environment in Rice - Fatima - 2022 - Journal of the Science of Food and Agriculture - Wiley Online Library
- Rising Carbon Dioxide and Global Nutrition: Evidence and Action Needed

References

Page 218

Hallgrave, D., Gilbert, S., et al. (2015) Regional dependencies in the beneficial effects of rising CO2 concentrations on crop water productivity. Nature Climate Change 5, 782–786. https://doi.org/10.1038/nclimate2665

Dr Delphine Devoy, lead author, IPCC WG1 WG2.

The data of the citation is incorrect, and I didn't find reference to this paper in the text of the chapter.

Return to table of contents

Supporting evidence

9.4 CO2 fertilisation and nutrient loss

Page 128

Optimal strategies will be location specific because they vary by crop, climate and soil type [the ref. 2015].

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

Developing new cultivars requires significant investments and generally take years per CO2AR groundwaters. Such germplasm will not be developed locally, nor will the necessary seeds for the use of optimal cultivar only applies to a specific location.

Return to table of contents

Supporting evidence

- CO2AR: A Model for Farmland's Complex Challenges

9.4 CO2 fertilisation and nutrient loss

Page 128

Second, identification of food products with micronutrients is already routine. Folic acid for B vitamin is added to flour and many other foods, iodine is added to table salt, and commercial breakfast cereals are fortified with iron and numerous vitamins. Thus, adding diverse supplements in the form of multivitamins tablets are inexpensive, widely available and routinely consumed.

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

Fortification and dietary supplements do not currently alleviate nutrient deficiencies; therefore, it is unlikely that such approaches will alleviate nutrient deficiencies in the future, particularly if they are not expanded to cover a greater range of micronutrients. 50% of women and girls worldwide (15–49 years) currently have iron deficiency anemia (IDA), and this is expected to rise to 60% by 2050 as a result of climate change. IDA is a factor in the CO2 stressed on 2 million people (Smith et al. 2017). Fortification and dietary supplement availability (McDougall 2017). The Dak CMS report ignores this and has not even considered the micronutrients that are commonly fortified in food and, such as lithium (for mental health), magnesium (for muscle and nerve function), blood pressure regulation, bone health and energy production) and iodine (Lobelde (2014); Dietary supplements do not address these critical contributors to micronutrient loss. Even if they did, as noted in the previous comment, supplements are not an adequate substitute for a nutrient dense diet. US Department of Agriculture and US Department of Health and Human Services (2010). Further, fortification of food or use of supplements will not address the health risk posed by increased toxic constituents in grain associated with elevated CO2 and CO2-induced warming.

Return to table of contents

Supporting evidence

- Anaemia Consortium Publications
- Prevalence of Iron Deficiency and Iron Deficiency Anemia in 5–6-Months-Aged U.S. Infants, 2005–2012 - Scott - 2016 - Nutrients - MDPI
- Iron Supplementation: What's New? - Blood - American Society of Hematology
- Hidden shift of the ionome of plants exposed to elevated CO2 depletes minerals at the base of human nutrition - eLife
- Food Composition Database Development Contributes to Micronutrient Studies in the United States, 2007–2016 - Journal of the American College of Nutrition: Vol 41, No 1

Back

9.4 CO2 fertilisation and nutrient loss

Page 128

Our concern about reliance on adaptive strategies is whether they are feasible in less developed countries. Micronutrient deficiency is already a problem in the developing world, and dietary supplements have proven to be an effective low-cost response [the ref. 2015].

Prof Kristie Ebi, professor of public health and climate change, Dr Becca Wasmund, associate professor of civil and environmental engineering.

As noted in the previous comment, micronutrient deficiency is a health problem in developing and developed countries; dietary supplements have not solved the problem. Micronutrient deficiency is a problem in the US (Bentley et al. 2017; Bentley et al. 2018; Smith and Myers (2019); Weyant et al. 2013; and Zhu et al. 2018). As supplements have been readily available, there is no evidence that they are economically infeasible. It is unlikely they will successfully ameliorate a future increased health burden caused by increased toxic constituents in grain. As noted in the previous comment, supplements will not address the health issues posed by increased toxic concentrations in grain expected with elevated CO2 and CO2-induced warming.

Return to table of contents

Supporting evidence

- Combining the effects of increased atmospheric carbon dioxide on protein, iron, and zinc availability and implications for global food supply - McDermid - 2019 - Global Change Biology - The Leading Publication for Global Health Implications of Nutrient Changes in Rice Grains: Risk Assessments Carbon Dioxide - Smith - 2019 - GeoHealth - Wiley Online Library
- Anthropogenic CO2 emissions and global food supply risk and vulnerability to artificial photosynthesis - A comparative modelling study - PLOS ONE.pdf
- Effects of elevated CO2 and temperature on the ionome, accumulation of essential nutrients and cadmium in wheat grown under current and future conditions - Journal of Experimental Botany

available that will need to be evaluated and adapted to local conditions.

Prof Kristie Ebi, professor of global health and Dr Becca Neumann, associate professor of civil and environmental engineering.

There is abundant evidence that under ideal conditions (i.e., adequate nutrients and water and optimal temperatures), crop productivity increases with rising CO2 levels (e.g., Ainsworth & Long 2020). These increases diminish or disappear when conditions are non-ideal (i.e., nutrients are limited, water and temperatures are sub-optimal). As CO2 concentrations increase, ambient temperatures will continue warming and weather patterns will shift, altering water and nutrient availability. It is not clear that boosts in productivity generated by elevated CO2 will compensate for the damage that crops particularly caused by these other stressors. In many parts of the world, even with adaptation, crop yields are expected to decline (Hultgren et al. (2021)). Further, discussed in the comments above, elevated CO2 is associated with declining nutrient density in crops. Therefore, it is disingenuous to state that there is "abundant evidence going back decades that rising CO2 levels benefit plants". This statement ignores the significant body of evidence host to plants associated with rising CO2 levels while ignoring the negative human health outcomes that will co-occur. The scientific consensus is that CO2-induced warming is harmful to US agriculture, leading to yield declines (Ebi et al. (2020), Hultgren et al. (2021); Lobell et al. (2011); Lobell and Field (2007); and Zhao et al. (2017)). Physiological studies of crops show there is an optimal temperature at which yields maximize. When temperatures are below or above this optimal point, yields decline.

Further, as outlined in the comments to Section 9.4 above, CO2-induced warming is associated with increased toxin levels in crops.It is factually inaccurate to state that "CO2 is not a pollutant with a set benefit to US agriculture". There are strategies, such as biofortification and supplementation, that can be employed to help mitigate nutrient dilution in crops (as discussed in Ebi et al. (2020). However, as outlined in the comments above, these approaches currently do not address nutrient deficiencies and therefore it is unrealistic to expect that they will be able to address future nutrient deficiencies exacerbated by rising CO2 levels and crop nutrient density. Further, these strategies will not solve the problem of increased concentrations of toxins in crops associated with increased CO2 and/or CO2-induced warming.

Quote given to Carbon Brief

Supporting evidence

1. Maize Cereal Grains Production and Use around the World | ACS Symposium Series
2. Impacts of climate change on global agriculture accounting for adaptation | Nature
3. Climate change impacts on crop yields: A review of empirical findings, statistical crop models, and machine learning methods | ScienceDirect
4. Climate Trends and Global Crop Production Since 1980 | Science
5. Effects of climate change on global agriculture: an interim assessment | SpringerLink
6. Temperature increase reduces global yields of major crops in four independent estimates | PNAS
7. Nutritional quality of crops in a high CO2 world: an agenda for research and technology development | IOPscience

10.5 Mortality from temperature extremes

Page 151

MISLEADING

Section 10.5.3

Dr Antonio Gasparrini, biostatistician and epidemiologist, London School of Hygiene & Tropical Medicine.

In relation to my area of research, the report cites two articles (Gasparrini (2015) and Zhao (2021)) in support of the statement that cold-related mortality far exceeds heat-related mortality in most of the regions. While true, this says little about the impact of *climate change* on temperature-related deaths. In fact, the focus should be on the respective (and opposite) changes in the heat and cold components, not their absolute values. In this respect, there is some evidence that the expected reduction in cold-related mortality will not offset the increase in heat-related mortality, in particular under more extreme climate change scenarios (Gasparrini et al. (2017); Masselot et al. (2022)). More importantly, the level of adaptation is hard to request to offset such a positive net effect should be very high, in the order of 90% reduction (Masselot (2025)).

Quote given to Carbon Brief

Supporting evidence

1. Mortality risk attributable to high and low ambient temperature: a multicountry observational study - PubMed
2. PDF
3. Temperature-related deaths and cold-related mortality under climate change: demographic and adaptation scenarios in 854 European cities - PubMed

11 Climate change, the economy, and the social cost of carbon

Page 156

FALSE

Ebi influential study in 2012 suggested that global warming would have growth in poor countries, but the finding has subsequently been found not to be robust. Studies that take full account of modelling uncertainties either find no evidence of a negative effect on global growth from CO2 emissions or find poor countries are likely to benefit as rich countries.

Prof Richard Tol, professor at the departments of economics, University of Sussex.

"The second sentence is wrong. The authors refer to "studies", but without references. Tol (2024) finds that the there-available studies jointly point to a negative impact of *climate change* on global economic growth. My less systematic reading of the literature since has not led me to change my mind. Their conclusion that '*poor countries*' are "likely to benefit" is again not backed up with references. Tol (2024), the only reference in the paragraph, concludes the opposite."

Quote from external source

Supporting evidence

1. A meta-analysis of the total economic impact of climate change - ScienceDirect

11.1 Climate change and economic growth

Page 157

MISLEADING

Not have past extreme weather events had a significant effect on US banks' performance (Bilicka et al. (2021)); would this even been shown to be beneficial for the finance and insurance sector (Mahadevan et al. (2025)).

Dr Kamiar Mohaddes, deputy director of the Cambridge executive MBA programme, University of Cambridge.

"The way our findings have been presented is misleading...Investigating the long-term macroeconomic effects of climate change across 40 US states, we provide evidence for the damage that climate change causes in the US using various economic indicators at the state level: growth rates of Gross State Product (GSP), GSP per capita (income), labour productivity (output per employee) as well as output growth in 10 economic sectors (e.g. agriculture, manufacturing, services, retail and wholesale trade). Just to be clear, (1) these are not 'small negative effects' and (2) we do indeed show that *climate change* has negative effects on income in the US. We show that while *heatwaves* shocks have level effects on (temporary growth impacts), *climate change* – by shifting the long-term average and variability of weather – impacts the US economy's ability to grow in the long term! We study economic activity in all sectors of the US economy –agriculture, forestry & fisheries; mining, construction, manufacturing, transport, communications and public utilities, wholesale trade, retail trade, financial services & property, services and government –and find that the impact of climate change on sectoral output growth is broad based – each of the 10 sectors considered is affected by at least one of the four climate variables."

Quote from external source

Supporting evidence

1. Climate change and economic activity: evidence from US states - Oxford Open Economics | Oxford Academic

11.1 Climate change and economic growth

Page 120

MISLEADING

Section 11.2 Models of the social cost of carbon

Dr Gernot Wagner, climate economist, Columbia Business School.

That section is based on a woefully outdated reading of the literature. In Moore et al (2024), we provide a detailed review of evidence of the latest such studies. We "perform a comprehensive synthesis of the evidence on the SCC, combining 1,823 estimates of the SCC from 147 studies with a survey of authors of these studies." In summary: "This synthesis elicited median values of $283 per tonne of CO2 for a 2020 pulse year (5% to 95% range: $32 to $874), higher than most official government estimates, including a 2021 update from the US EPA." So far from there being 'evidence for low SCC' (DOE report p.125), sadly, there is ample evidence of the opposite.

Quote given to Carbon Brief

Supporting evidence

1. Synthesis of evidence yields high social cost of carbon due to structural model variation and uncertainties | PNAS

11.2 Models of the social cost of carbon

Page 121

MISLEADING

Economists use IAMs to compute the SCC. Two of the best known are the Climate Framework for Uncertainty, Negotiation and Distribution ("FUND", Tol 1997) and Nordhaus' DICE. EPA (2023) introduced new uses for its recent work. IAMs embed a "damage function" or set of functions relating ambient temperature to local economic conditions. The assumptions embedded in the damage function will largely determine the resulting SCC. IAMs also assume a long-term discount rate or, as in DICE, compute the optimal internal discount rate as part of the solution.

Prof Richard Tol, professor at the departments of economics, University of Sussex.

"The literature is vast. I counted 446 papers with estimates. There are numerous circumstances, and two handfuls of meta-analyses (e.g. Tol (2023) and Moore et al. (2024)). Instead, the authors point out two reviews, which omits the most influential papers and misses key insights. Cherry-picking may be a better term than review."

Quote from external source

Supporting evidence

1. Social cost of carbon estimates have increased over time | Nature Climate Change
2. Synthesis of evidence yields high social cost of carbon due to structural model variation and uncertainties | PNAS

11.2 Models of the social cost of carbon

Page 123

FALSE

Tol (2017) estimates that the private benefit of carbon is large relative to the social cost.

Prof Richard Tol, professor at the departments of economics, University of Sussex.

"This paper was never published in a peer-reviewed journal and is therefore not admissible by the rules of the US government. The paper was peer-reviewed and rejected, because my private benefit jof carbon] is an average, whereas the social cost is a marginal. The two cannot be compared (unless you make a ridiculous assumption about linearity). I still hope to fix the paper one day. As it stands, however, the comparison is wrong."

Quote from external source

Supporting evidence

1. The Private Benefit of Carbon and its Social Cost

11.2 Models of the social cost of carbon

Page 125

MISLEADING

Dietz et al, incorporated several potential tipping points (abrupt changes) into an SCC model and found they added about 25% to the estimate, mainly associated with thawing permafrost and release of methane hydrates. However, the IPCC considers this scenario very unlikely (AR6 Technical Summary p. 109).

Dr Gernot Wagner, climate economist, Columbia Business School.

Dietz et al., in this main specification, finds a median increase in the SCC of 'about 25%' (27%) and a mean increase of over 40% (42%). That comparison of median and mean points to a heavily right-skewed distribution. Figure 2 shows as much. It also shows, for example, that we find a ~10% chance that these tipping points alone increase the SCC by at least 100% – that is, double it.

Quote given to Carbon Brief

Supporting evidence

1. Economic impacts of tipping points in the climate system | PNAS

Glossary

IPCC: The Intergovernmental Panel on Climate Change, the UN body for assessing the science related to climate change. IPCC works in five-to-eight-year cycles and comprises three working groups of scientists, each focusing on different aspects of climate science and the response to climate change. Each assessment cycle, scientists appointed by the IPCC produce three reports summarising the latest climate science, as well as "special reports" and a synthesis report, which serve as key inputs to international climate negotiations.

WG1/2/3: The three working groups of the IPCC. Working Group I (WG1) deals with the physical science basis of climate change, Working Group II (WG2) with climate change impacts, adaptation and vulnerability and Working Group III (WG3) with mitigation of climate change.

AR6: The sixth assessment report of the IPCC, which was published in 2013-14.

AR5: The fifth assessment report of the IPCC, which was published in 2021-23.

CMIP5/6: The Coupled Model Intercomparison Project (CMIP) is the international modelling effort that feeds into the IPCC's influential assessments. CMIP5 was developed for AR5, while CMIP6 was developed for AR6.

Methodology

The DoB report references around 550 sources. Of these, around 240 are peer-reviewed research papers. Other sources include books, commentary articles and letters published in journals, as well as "gray" literature, such as study preprints, blog posts, news articles and government websites.

For this factcheck, Carbon Brief emailed the corresponding authors of all 240 peer-reviewed research papers referenced in the preceding author did not respond or the email address bounced, Carbon Brief contacted another report author – typically the lead author, if separate to the corresponding author.

In 20 cases, the corresponding author was also an author on the DoE report itself. These authors were not invited to factcheck their own work.

Carbon Brief asked the corresponding authors to read the section of the report that cites their work and to identify if it uses their work to make "misleading" or "false" statements.

False was described as "statements/statistics that are factually incorrect or untrue".

Misleading was described as "statements/statistics that are presented in a way that distort the meaning, create a false impression or omit/cherry-pick crucial details".

(Carbon Brief also invited various topic experts to contribute their responses. Respondents were also invited to forward Carbon Brief's request to other experts.)

Carbon Brief's factcheck also includes comments from scientists who have already reacted to the DoE report in public, via media articles, social media or blog posts. Links to these sources are included with the relevant responses.

Carbon Brief ensured that any scientists contributing to the factcheck in an anonymous capacity have the appropriate expertise to comment on the sections of the report that they contributed.

Any pages with "misleading" and "false" statements are shown in orange and red, respectively, in the main graphic. Pages that are coloured are either accurate, or have not been factchecked.

There has been some speculation that artificial-intelligence tools might have been used to generate some parts of the report, particularly the citations, due to some "hallucinations" – type mistakes. For example, a Carbon Brief article from 2017 is cited, but the credited author is mistakenly listed as another Carbon Brief journalist.

However, Carbon Brief can samples of the report's main text through several AI text checker tools and found there is a "very low probability" they were generated using AI. It was not possible to test the citations, due to the paywalls.

Update: This article was updated once on 22.08.25 to make further errors in the report to email info@carbonbrief.org.